

Rational homology and homotopy of high dimensional string links

Paul Arnaud Songhafou Tsopméné
Victor Turchin

Abstract

Arone and Turchin obtained the Hodge decomposition in the homology and homotopy of the high dimensional analogues of spaces of long knots. Next they showed that when the dimensions are in the stable range, the rational homology and homotopy of these latter spaces can be calculated as the homology of a direct sum of certain finite graph-complexes that they described explicitly. In this paper, we generalize all these results to high dimensional analogues of spaces of string links.

The methods of our paper are applicable in the range when the ambient dimension is at least twice the maximal dimension of a link component plus two, which in particular guarantees that the spaces under the study are connected. However, we conjecture that our homotopy graph-complex computes the rational homotopy groups of links spaces always when codimension is greater than two, i.e. always when the Goodwillie-Weiss calculus is applicable. Using Haefliger's approach to study isotopy classes of higher dimensional links, we confirm our conjecture at the level of π_0 .

0 Introduction

In this paper we further develop the connection between the study of embedding spaces, operad theory, and graph-complexes. Our results are partially influenced by Bar Natan's work [4], where he describes the space dual to weight systems (counting Vassiliev invariants) of string links in $\mathbb{R}^3$ as a certain space of univalent graphs modulo AS and IHX relations, with univalent vertices being colored by the components of the links, see also [7, Section 5.10] for more details. This space is built from the components of the modular envelop $\mathbf{Mod}(\text{Lie})$ of the operad of Lie algebras [25]. One of our main results says that the rational homotopy of the space of higher-dimensional string links is described as the homology of a graph-complex similarly built from the components of the modular envelop of L_∞ .¹ In particular Bar Natan's space of univalent graphs with colored univalent vertices appears as the bottom degree homology of our complexes. In this paper we produce two different graph-complexes computing the rational homotopy of spaces of string links. One of them is built from the components of $\mathbf{Mod}(L_\infty)$ and another one has building blocks described in terms of the homology of configuration spaces of points in Euclidean spaces. Both complexes resemble those appearing in the Bott-Taubes type integration, see for example [37, 38]. The obtained complexes all split into a direct sum of finite complexes. In the follow up paper [41], we compute the generating function of the Euler characteristics of the summands of these complexes (both for the homology and homotopy). As a byproduct result in [41], we compute the supercharacter of the symmetric group action on the positive arity components of $\mathbf{Mod}(L_\infty)$.

¹For the notion of a modular operad and modular envelop, sometimes called modular closure, see [18, 31, 25].

Let d be the dimension of the ambient space. Let $r \geq 1$, and let $m_1, \dots, m_r \geq 1$ be r integers. A *high-dimensional string link of r strands* is a smooth embedding $f: \coprod_{i=1}^r \mathbb{R}^{m_i} \hookrightarrow \mathbb{R}^d$ that coincides outside a compact set with a fixed embedding $\iota: \coprod_{i=1}^r \mathbb{R}^{m_i} \hookrightarrow \mathbb{R}^d$ affine on each component. When r is understood we will just say *high-dimensional string link*. The space of high-dimensional string links, denoted $\text{Emb}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$, is the collection of all high-dimensional string links endowed with the weak $\mathcal{C}^\infty$ -topology. Similarly, we define the space $\text{Imm}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$ of smooth immersions $\coprod_{i=1}^r \mathbb{R}^{m_i} \hookrightarrow \mathbb{R}^d$ that coincide outside a compact set with ι . By definition, there is an inclusion $\text{Emb}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) \hookrightarrow \text{Imm}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$, and its homotopy fiber over ι , denoted $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$, is the space under our study in this paper. Notice that the case $r = 1$ was extensively studied in [2] and [3] by Arone and Turchin. They obtain the Hodge decomposition in the rational homology $H_*(\overline{\text{Emb}}_c(\mathbb{R}^m, \mathbb{R}^d), \mathbb{Q})$ and in the rational homotopy $\mathbb{Q} \otimes \pi_* \overline{\text{Emb}}_c(\mathbb{R}^m, \mathbb{R}^d)$, $d > 2m + 1$. Then they explicitly describe graph-complexes computing $H_*(\overline{\text{Emb}}_c(\mathbb{R}^m, \mathbb{R}^d), \mathbb{Q})$ and $\mathbb{Q} \otimes \pi_* \overline{\text{Emb}}_c(\mathbb{R}^m, \mathbb{R}^d)$.

Because of the high codimension, the space $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$ is connected and moreover its homotopy type does not depend on the choice of the fixed affine embedding ι . In particular we can choose ι in a way that all connected components of the fixed unlink are parallel to a given line. Concatenation of links along the direction of this line defines an H -space structure on $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$, which implies that the rational homology of $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$ is a cofree cocommutative coalgebra cogenerated by the rational homotopy. The same is true for the graph-complexes that we study.

In order to describe our graph-complexes, we need Theorem 0.1 below, which is a generalization of [2, Theorem 1.3]. Before stating Theorem 0.1, we will recall two notions. The first one is that of right Ω -modules. Let Ω be the category of finite unpointed sets whose morphisms are surjections. A *right Ω -module* is a contravariant functor from Ω to any given category. We denote by Rmod_Ω the category of right Ω -modules of chain complexes over $\mathbb{Q}$. Abusing notation let $i \in \Omega$ denote the set $\{1, \dots, i\}$. The category of right Ω -modules of $\mathbb{Q}$ -vector spaces is an abelian category with the set of projective generators $\mathbb{Q}[\text{hom}_\Omega(i, -)]$, $i \geq 0$. Using this fact we endow Rmod_Ω with the structure of a (cofibrantly generated) model category, whose weak equivalences are quasi-isomorphisms, and fibrations are degreewise surjective maps [26]. (We call it projective model structure.) We denote by $\text{Rmod}_\Omega(-, -)$ the space (chain complex) of maps between two Ω -modules, and by $\text{hRmod}_\Omega(-, -)$ the derived mapping space. For specific computations we will need to apply this construction only to Ω -modules with zero differential, in which case $\text{hRmod}_\Omega(-, -)$ can be expressed as a product of certain Ext groups.

As an example of a right Ω -module (the one that appears in Theorem 0.1), we have the following. Let X be a pointed topological space. Define the functor $X^{\wedge \bullet}$ that assigns to the set $\{1, 2, \dots, k\}$ the space $X^{\wedge k} = \underbrace{X \wedge \dots \wedge X}_k$ (here " $\wedge$ " is the smash product operation). By $X^{\wedge 0}$ we mean the two-point space – zero dimensional pointed sphere S^0 . The morphisms are induced by the diagonal maps. This functor is a right Ω -module with values in topological spaces. Thus the functors $\tilde{C}_*(X^{\wedge \bullet})$ (here $\tilde{C}_*(-)$ denotes the reduced chain complex functor), $\tilde{H}_*(X^{\wedge \bullet})$ (with zero differential) are objects of Rmod_Ω .

The second notion we need to recall is that of right Γ -modules, which is the same as the previous one with the category Ω replaced by the category Γ . Recall that objects in Γ are finite pointed sets, while morphisms are pointed maps. As before we have the category Rmod_Γ (of right Γ -modules in chain complexes). We similarly denote by $\text{Rmod}_\Gamma(-, -)$ the mapping space and by $\text{hRmod}_\Gamma(-, -)$ the derived mapping space between two right Γ -modules. The example of a right Γ -module we look at is $H_*(C(\bullet, \mathbb{R}^d), \mathbb{Q})$, $d \geq 2$, where $C(k, \mathbb{R}^d)$ denotes the configuration space of k labeled points in $\mathbb{R}^d$. A way to see that $H_*(C(\bullet, \mathbb{R}^d), \mathbb{Q})$ is a right

Γ -module is the fact that it is an infinitesimal bimodule (see [2, Definition 3.8] or [42, Definition 4.1] for the definition of an *infinitesimal bimodule*) over $\text{Com} = H_0(C(\bullet, \mathbb{R}^d), \mathbb{Q})$ since there is an obvious morphism of operads $\text{Com} \longrightarrow H_*(C(\bullet, \mathbb{R}^d), \mathbb{Q})$, the target $H_*(C(\bullet, \mathbb{R}^d), \mathbb{Q}) = H_*(\mathcal{B}_d(\bullet), \mathbb{Q})$ is the homology of the little discs operad $\mathcal{B}_d(\bullet)$, which is well known to be the graded Poisson algebras operad [8]. It is also well known that an infinitesimal bimodule over Com is the same thing as a right Γ -module (see [2, Corollary 4.10] or [42, Lemma 4.3]). One can also show that the sequence $\mathbb{Q} \otimes \pi_* C(\bullet, \mathbb{R}^d)$, $d \geq 3$, has a natural structure of a right Γ -module.

In [35] Pirashvili constructs a functor (called *cross effect*) $\text{cr}: \text{Rmod}_\Gamma \longrightarrow \text{Rmod}_\Omega$ that turns out to be an equivalence of categories. Let $\hat{H}_*(C(\bullet, \mathbb{R}^d), \mathbb{Q})$ (respectively $\mathbb{Q} \otimes \hat{\pi}_* C(\bullet, \mathbb{R}^d)$) denote the cross effect of $H_*(C(\bullet, \mathbb{R}^d), \mathbb{Q})$ (respectively the cross effect of $\mathbb{Q} \otimes \pi_* C(\bullet, \mathbb{R}^d)$).

We are now ready to state our first result, which concerns the rational homology of $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$.

Theorem 0.1. *For $d > 2\max\{m_i \mid 1 \leq i \leq r\} + 1$, there is an isomorphism*

$$H_*(\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \mathbb{Q}) \cong H\left(\text{hRmod}_\Omega\left(\tilde{H}_*((\vee_{i=1}^r S^{m_i})^{\wedge \bullet}, \mathbb{Q}), \hat{H}_*(C(\bullet, \mathbb{R}^d), \mathbb{Q})\right)\right). \quad (0.1)$$

Our second result concerns the rational homotopy of $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$, and is the following.

Theorem 0.2. *For $d > 2\max\{m_i \mid 1 \leq i \leq r\} + 1$, there is an isomorphism*

$$\mathbb{Q} \otimes \pi_* \overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) \cong H\left(\text{hRmod}_\Omega\left(\tilde{H}_*((\vee_{i=1}^r S^{m_i})^{\wedge \bullet}, \mathbb{Q}), \mathbb{Q} \otimes \hat{\pi}_* C(\bullet, \mathbb{R}^d)\right)\right). \quad (0.2)$$

The main ingredients in the proof of Theorem 0.2 are Theorem 0.1 and hairy graph-complexes that will be explicitly described in Subsection 2.1.

In the following lines, we will see the Hodge decomposition in the homology $H_*(\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \mathbb{Q})$. Notice that our decomposition can be viewed as a generalization of a result of Pirashvili [36], which concerns the Hodge decomposition for the S^m -homology of any right Γ -module (a.k.a. Hodge decomposition for the higher order Hochschild homology). We say "generalization" because, as in [40] or as we will see in Section 1, it is not difficult to show that the homology $H_*(\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \mathbb{Q})$ is the $\vee_{i=1}^r S^{m_i}$ -homology of the Γ -module $H_*(C(\bullet, \mathbb{Q}))$. The Hodge-type splitting that we study can be in the same way defined for the $\vee_{i=1}^r S^{m_i}$ higher Hochschild homology of any Γ -module and appears because $C_*((\vee_{i=1}^r S^{m_i})^{\times \bullet}, \mathbb{Q})$ is formal as a right Γ -module (which itself is equivalent to the fact that its cross effect Ω -module $\tilde{C}_*((\vee_{i=1}^r S^{m_i})^{\wedge \bullet}, \mathbb{Q})$ is formal). More generally such splitting in higher Hochschild homology takes place for any suspension and has been recently studied in [45].

The right hand sides of (0.1) and (0.2) can be split as follows. For any homomorphism of finite groups $H \rightarrow G$, let Ind_H^G denote the induction functor from H -representations to G -representations, which is left adjoint to the restriction functor.² In most of the cases G will be a symmetric group and H be its subgroup. Let us consider the right Ω -module $\tilde{H}_*((\vee_{i=1}^r S^{m_i})^{\wedge \bullet}, \mathbb{Q})$ that appears in (0.1). By definition, its k th component is

²Explicitly, since we deal with the right action: $\text{Ind}_H^G V = V \otimes_{\mathbb{Q}[H]} \mathbb{Q}[G]$.

given by

$$\begin{aligned}
\tilde{H}_*((\bigvee_{i=1}^r S^{m_i})^{\wedge k}, \mathbb{Q}) &= \tilde{H}_*\left(\underbrace{(\bigvee_{i=1}^r S^{m_i}) \wedge \cdots \wedge (\bigvee_{i=1}^r S^{m_i})}_k, \mathbb{Q}\right) \\
&\cong \tilde{H}_*\left(\bigvee_{i_1, \dots, i_k=1}^r (S^{m_{i_1}} \wedge \cdots \wedge S^{m_{i_k}}), \mathbb{Q}\right) \\
&\cong \tilde{H}_*\left(\bigvee_{s_1 + \cdots + s_r = k} \bigvee_{\frac{k!}{s_1! \cdots s_r!}} S^{s_1 m_1} \wedge \cdots \wedge S^{s_r m_r}, \mathbb{Q}\right).
\end{aligned}$$

Thus

$$\tilde{H}_*((\bigvee_{i=1}^r S^{m_i})^{\wedge k}, \mathbb{Q}) \cong_{\Sigma_k} \bigoplus_{s_1 + \cdots + s_r = k} \text{Ind}_{\Sigma_{s_1} \times \cdots \times \Sigma_{s_r}}^{\Sigma_k} \tilde{H}_*(S^{s_1 m_1} + \cdots + S^{s_r m_r}, \mathbb{Q}). \quad (0.3)$$

This implies the following isomorphism of right Ω -modules

$$\tilde{H}_*((\bigvee_{i=1}^r S^{m_i})^{\wedge \bullet}, \mathbb{Q}) \cong \bigoplus_{s_1, \dots, s_r \geq 0} Q_{s_1 \cdots s_r}^{m_1 \cdots m_r}, \quad (0.4)$$

where $Q_{s_1 \cdots s_r}^{m_1 \cdots m_r}$ is the right Ω -module defined by

$$Q_{s_1 \cdots s_r}^{m_1 \cdots m_r}(k) = \begin{cases} 0 & \text{if } k \neq s_1 + \cdots + s_r; \\ \text{Ind}_{\Sigma_{s_1} \times \cdots \times \Sigma_{s_r}}^{\Sigma_k} \tilde{H}_*(S^{s_1 m_1} + \cdots + S^{s_r m_r}, \mathbb{Q}) & \text{if } k = s_1 + \cdots + s_r. \end{cases} \quad (0.5)$$

Any non-bijective surjection acts on $Q_{s_1 \cdots s_r}^{m_1 \cdots m_r}$ automatically as zero.

On the other hand, we have the splittings

$$\hat{H}_*(C(\bullet, \mathbb{R}^d), \mathbb{Q}) = \prod_{t \geq 0} \hat{H}_{t(d-1)}(C(\bullet, \mathbb{R}^d), \mathbb{Q}). \quad (0.6)$$

$$\mathbb{Q} \otimes \hat{\pi}_* C(\bullet, \mathbb{R}^d) = \prod_{t \geq 0} \mathbb{Q} \otimes \hat{\pi}_{t(d-2)+1} C(\bullet, \mathbb{R}^d). \quad (0.7)$$

Combining (0.1), (0.2), (0.4), (0.6), and (0.7), it is not difficult to get the following splittings

$$\begin{aligned}
H_*(\overline{\text{Emb}}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \mathbb{Q}) &\cong \prod_{s_1, \dots, s_r, t} H_* \text{hRmod}_{\Omega} \left(Q_{s_1 \cdots s_r}^{m_1 \cdots m_r}, \hat{H}_{t(d-1)}(C(\bullet, \mathbb{R}^d), \mathbb{Q}) \right) \\
&\cong \bigoplus_{s_1, \dots, s_r, t} H_* \text{hRmod}_{\Omega} \left(Q_{s_1 \cdots s_r}^{m_1 \cdots m_r}, \hat{H}_{t(d-1)}(C(\bullet, \mathbb{R}^d), \mathbb{Q}) \right). \quad (0.8)
\end{aligned}$$

$$\begin{aligned} \mathbb{Q} \otimes \pi_* (\overline{\text{Emb}}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)) &\cong \prod_{s_1, \dots, s_r, t} H_* \text{hRmod}_{\Omega} (Q_{s_1 \dots s_r}^{m_1 \dots m_r}, \mathbb{Q} \otimes \widehat{\pi}_{t(d-2)+1} C(\bullet, \mathbb{R}^d)) \\ &\cong \bigoplus_{s_1, \dots, s_r, t} H_* \text{hRmod}_{\Omega} (Q_{s_1 \dots s_r}^{m_1 \dots m_r}, \mathbb{Q} \otimes \widehat{\pi}_{t(d-2)+1} C(\bullet, \mathbb{R}^d)). \end{aligned} \quad (0.9)$$

The product is replaced by the direct sum because only finitely many factors contribute for any given degree. We use here that $d > 2\max\{m_i \mid 1 \leq i \leq r\} + 1$. This can be seen from the explicit description of our complexes analogously to the case $r = 1$, see Remark 2.4. (For (0.9), this is true even for a weaker constraint $d > \max\{m_i \mid 1 \leq i \leq r\} + 2$.) The splitting by $s_1, \dots, s_r$ will be called Hodge splitting and the splitting by t will be called splitting by complexity.

The current paper is the first one of our project. In the second one [41], we determine the generating functions of the Euler characteristics of the summands in (0.8) and (0.9). By the time when our project was completed, another work by Fresse, Willwacher and the second author appeared [16], where among other things, by very different techniques, the statement of Theorem 0.2, equivalently (0.9), was improved for $r = 1$ to codimension $d - m_1 > 2$.³ Using the classical study of the isotopy classes of higher dimensional spherical links due to A. Haefliger and others [24, 10], we confirmed, see Theorem 3.2, that (0.9) holds at the level of π_0 always when codimension > 2 . Based on these facts, we conjecture that this identity (0.9) is true for $* \geq 0$ and $d - \max\{m_i \mid 1 \leq i \leq r\} > 2$, see Conjecture 3.1.

Outline of the paper.

- In Section 1 we prove Theorem 0.1, which determines the rational homology of the space of high dimensional string links. The proof is essentially based on a version, developed by Arone and Turchin in [2], of the Goodwillie-Weiss embedding calculus [20, 46]. It uses the relative formality theorem (of the inclusion $\mathcal{B}_n \hookrightarrow \mathcal{B}_d$ of the little n -disks operad inside the little d -disks operad) proved for $d \geq 2n + 1$ in [28] and improved to the range $d \geq n + 2$ in [44]. It also uses the fact that the right Ω -module $\tilde{C}_*((\bigvee_{i=1}^r S^{m_i})^{\wedge \bullet}, \mathbb{Q})$ is formal in the category Rmod_{Ω} .
- In Section 2 we prove Theorem 0.2, which determines the rational homotopy of $\overline{\text{Emb}}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$. The plan of the proof is the same as that of [3, Theorem 1.1 (ii)], for which our theorem is a generalization. In order to prove [3, Theorem 1.1 (ii)], the authors, Arone and Turchin, explicitly describe *hairy graph-complexes* computing the rational homology and the rational homotopy of the space $\overline{\text{Emb}}_c(\mathbb{R}^m, \mathbb{R}^d)$. Our complexes, explicitly described in Section 2.1, are the same as theirs except that the external vertices of our graphs are colored by the components of the links. They are obtained by taking the injective resolution of the target Ω -modules in (0.8) and (0.9). Theorem 0.2 follows from the fact that the homotopy hairy graph-complex turns out to be exactly the subcomplex of connected graphs of the homology hairy graph-complex.
- Section 3 is devoted to Conjecture 3.1 saying that our hairy graph-complex computes the rational homotopy of $\overline{\text{Emb}}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$ for any codimension > 2 . We show Theorem 3.2 that the statement of the conjecture is true at the level of π_0 . From the point of view of the hairy graph-complex only its

³More precisely they proved that the rational homotopy groups of the little discs operads mapping space $\text{hOper}(\mathcal{B}_m, \mathcal{B}_d)$, $d - m > 2$, are computed by the (shifted) hairy graph-complex $\mathcal{E}_{\pi}^{m,d}$ that describes explicitly the right-hand side of (0.9) (see Section 2). (In [16] this complex is denoted by $\text{HGC}_{m,d}$.) The latter result implies ours thanks to the delooping result $\overline{\text{Emb}}_c(\mathbb{R}^m, \mathbb{R}^d) \simeq \Omega^{m+1} \text{hOper}(\mathcal{B}_m, \mathcal{B}_d)$ from [5], see also [11, 12, 14, 43].

tree part can contribute to degree zero. Another way to say it is that the rational additive invariants of higher dimensional string links can all be expressed as generalized Milnor type invariants [6, 21, 27, 32].⁴

- In Section 4 we explicitly describe graph-complexes computing the rational (co)homology and homotopy of $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$, which are obtained by taking projective resolution of the source Ω -modules in (0.8) and (0.9). We use here the fact that a right Ω -module is the same thing as a right module over the commutative non-unital operad Com_+ and also the Koszul duality between Com_+ and the operad Lie of Lie algebras. The obtained complexes are thus called *Koszul complexes*. We use them in the follow up paper [41] for the computations of the Euler characteristics of the summands in (0.8) and (0.9). We also show that natural truncations of the Koszul complexes can be used to determine the stages of the Goodwillies-Weiss towers $T_\bullet C_*(\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \mathbb{Q})$ and (conjecturally) $\mathbb{Q} \otimes \pi_* T_\bullet \overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$.

Acknowledgements This work has been supported by Fonds de la Recherche Scientifique-FNRS (F.R.S.-FNRS), that the authors acknowledge. It has been also supported by the Kansas State University (KSU), where this paper was partially written during the stay of the first author, and which he thanks for hospitality.

1 The rational homology of $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$

The goal of this section is to prove Theorem 0.1 announced in the introduction.

The first and main step in proving Theorem 0.1 is Theorem 1.4. Propositions 1.2 and 1.3 below will be used in the proof of the latter theorem. Define $m = \max\{m_i \mid 1 \leq i \leq r\}$. Without changing the homotopy type of $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$ we can assume that the fixed affine embedding $\iota: \coprod_{i=1}^r \mathbb{R}^{m_i} \hookrightarrow \mathbb{R}^d$ can be factored through $\mathbb{R}^{m+1}$. Moreover let us assume that ι restricted on each component $\mathbb{R}^{m_i}$ can be factored through an affine m -space $V_i \subset \mathbb{R}^{m+1}$, so that all V_i , $i = 1 \dots r$, are parallel to each other and disjoint. For a space X let $\widehat{X}$ be its one point compactification. Define M_i as a tubular neighborhood of $\widehat{\iota(\mathbb{R}^{m_i})}$ inside $\widehat{V_i}$ minus the point at infinity. Simialrly define M as an $(m+1)$ -submanifold of $\mathbb{R}^{m+1}$ obtained as a tubular neighborhood of the wedge of spheres $\widehat{\iota(\coprod_{i=1}^r \mathbb{R}^{m_i})}$ in $\widehat{\mathbb{R}^{m+1}}$ minus ∞ .

Remark 1.1. *We don't need it for any of our arguments, but one can show that each M_i is diffeomorphic to $\mathbb{R}^m \setminus S^{m-m_i-1}$, where S^{m-m_i-1} is an unknotted $(m-m_i-1)$ -sphere in $\mathbb{R}^m$. (By (-1) -sphere as usual we understand the empty set.) One can also show that M is diffeomorphic to $\mathbb{R}^{m+1} \setminus \bigvee_{i=1}^r S^{m-m_i}$, where $\bigvee_{i=1}^r S^{m-m_i}$ is an unknotted wedge of spheres in $\mathbb{R}^{m+1}$.*

Proposition 1.2. *The natural restriction maps*

$$\overline{\text{Emb}}_c(M, \mathbb{R}^d) \xrightarrow{\simeq} \overline{\text{Emb}}_c(\coprod_{i=1}^r M_i, \mathbb{R}^d) \xrightarrow{\simeq} \overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) \quad (1.1)$$

are weak equivalences.

Proof. It is easy to see that the embedding spaces $\text{Emb}_c(M, \mathbb{R}^d)$ and $\text{Emb}_c(\coprod_{i=1}^r M_i, \mathbb{R}^d)$ (respectively immersion spaces $\text{Imm}_c(M, \mathbb{R}^d)$ and $\text{Imm}_c(\coprod_{i=1}^r M_i, \mathbb{R}^d)$) are homotopy equivalent to the spaces of partially framed long embeddings $\coprod_{i=1}^r \mathbb{R}^{m_i} \hookrightarrow \mathbb{R}^d$ (respectively immersions $\coprod_{i=1}^r \mathbb{R}^{m_i} \looparrowright \mathbb{R}^d$). By a “partially framing” of a long link we mean that the i th component of a link has $(m+1-m_i)$ in case of M and $(m-m_i)$ in

⁴We also believe that such invariants can be expressed as Bott-Taubes type integrals similarly to the Haefliger invariant [37].

case of $\coprod_{i=1}^r M_i$ everywhere linearly independent sections of the normal bundle, which must coincide with a fixed constant partial framing outside a compact set. When we pass to the homotopy fiber or in other words look at the spaces of partially framed embeddings modulo partially framed immersions this partial framing cancels out. $\square$

One of the reasons we need all the components of our links to be of the same dimension is that we want to apply the theory [46, 20] of Goodwillie-Weiss manifold functor calculus. If F is a functor from a category of open subsets of a manifold (which serves as an input in that theory), the k th approximation of F is denoted $T_k F$, $k \leq \infty$. The sequence $\{T_k F\}_{0 \leq k \leq \infty}$ is usually called the *Taylor tower* associated to F , which converges, under good assumptions, to F . In our situation we apply the functor calculus to the categories of open sets of M , respectively of $\coprod_{i=1}^r M_i$, whose complement in $\mathbb{R}^{m+1}$, respectively in $\coprod_{i=1}^r \mathbb{R}^m$, is compact.

Proposition 1.3. *The restriction map $\overline{\text{Emb}}_c(M, \mathbb{R}^d) \xrightarrow{\simeq} \overline{\text{Emb}}_c(\coprod_{i=1}^r M_i, \mathbb{R}^d)$ induces an equivalence of Goodwillie-Weiss towers*

$$T_k C_* \overline{\text{Emb}}_c(M, \mathbb{R}^d) \xrightarrow{\simeq} T_k C_* \overline{\text{Emb}}_c\left(\coprod_{i=1}^r M_i, \mathbb{R}^d\right), \quad 0 \leq k \leq \infty.$$

Proof. For $k < \infty$, both $T_k C_* \overline{\text{Emb}}_c(M, \mathbb{R}^d)$ and $T_k C_* \overline{\text{Emb}}_c(\coprod_{i=1}^r M_i, \mathbb{R}^d)$ are obtained as a homotopy limit over a finite diagram encoding the handle attachments and iteration of the limits over corresponding $(k+1)$ -subcubical diagrams. One has a natural transformation induced by restriction between the two diagrams which is an objectwise equivalence. As a consequence the homotopy limits must be also equivalent. The case $k = \infty$ is obtained by taking $k \rightarrow \infty$. $\square$

Recalling that if X is a pointed topological space, then the sequence $X^{\times \bullet} = \underbrace{\{X \times \cdots \times X\}}_k_{k \geq 0}$ is equipped with a natural structure of a right Γ -module: to every finite pointed set we assign the space of base preserving maps from it to X . We have the following theorem in which the notations " hRmod_Γ " and " $C(\bullet, \mathbb{R}^d)$ " were introduced in Section 0.

Theorem 1.4. *For $d > 2\max\{m_i \mid 1 \leq i \leq r\} + 1$, there is a quasi-isomorphism*

$$C_*(\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)) \otimes \mathbb{Q} \simeq \text{hRmod}_\Gamma(C_*((\bigvee_{i=1}^r S^{m_i})^{\times \bullet}, \mathbb{Q}), H_*(C(\bullet, \mathbb{R}^d), \mathbb{Q})).$$

Proof. Let us start with the following weak equivalences.

$$\begin{aligned} C_* \overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) &\simeq C_* \overline{\text{Emb}}_c(\coprod_{i=1}^r M_i, \mathbb{R}^d) && \text{by Proposition 1.2} \\ &\simeq T_\infty C_* \overline{\text{Emb}}_c(\coprod_{i=1}^r M_i, \mathbb{R}^d) && \\ &\simeq T_\infty C_* \overline{\text{Emb}}_c(M, \mathbb{R}^d) && \text{by Proposition 1.3.} \end{aligned} \tag{1.2}$$

The second equivalence is the fact that the Goodwillie-Weiss tower for singular chains of embeddings converges for $d > 2m + 1$ [46].

By the Arone-Turchin work [2], one has [40, Proposition 2.11], which states that if $N \subseteq \mathbb{R}^n$ is the complement of a compact subset of $\mathbb{R}^n$, then

$$T_\infty C_*(\overline{\text{Emb}}_c(N, \mathbb{R}^d), \mathbb{Q}) \simeq \text{hRmod}_\Gamma(C_*(\widehat{N}^{\times \bullet}), H_*(C(\bullet, \mathbb{R}^d); \mathbb{Q})), \tag{1.3}$$

where $\widehat{N}$ is an open subset of the one-point compactification of $\mathbb{R}^n$ consisting of N and ∞ . This space is pointed at ∞ . In [40, Proposition 2.11] this result is stated in the range $d > 2n + 1$ as it uses the relative formality of the little discs operads which was proved at that time only in this range [28]. As it turned the relative formality holds in larger range $d > n + 1$ [44] implying (1.3) for $d > n + 1$.

By rewriting (1.3) with our manifold $M \subseteq \mathbb{R}^{m+1}$, we obtain the following weak equivalence (for $d > m + 2$):

$$T_\infty C_*(\overline{\text{Emb}}_c(M, \mathbb{R}^d), \mathbb{Q}) \simeq \text{hRmod}_{\Gamma}(C_*(\widehat{M}^{\times \bullet}), H_*(C(\bullet, \mathbb{R}^d); \mathbb{Q})). \quad (1.4)$$

Combining now (1.2) and (1.4), and using the fact that $\widehat{M}$ is homotopy equivalent to $\bigvee_{i=1}^r S^{m_i}$ we get exactly the statement of our theorem. $\square$

In [35] Pirashvili constructs a functor (called the *cross effect*)

$$cr: \text{Rmod}_{\Gamma} \longrightarrow \text{Rmod}_{\Omega} \quad (1.5)$$

that turns out to be an equivalence of categories. Define

$$\widehat{H}_*(C(\bullet, \mathbb{R}^d), \mathbb{Q}) = cr(H_*(C(\bullet, \mathbb{R}^d), \mathbb{Q})).$$

By noticing that the cross effect of $C_*((\bigvee_{i=1}^r S^{m_i})^{\times \bullet}, \mathbb{Q})$ is $\widetilde{C}_*((\bigvee_{i=1}^r S^{m_i})^{\wedge \bullet}, \mathbb{Q})$, where $\widetilde{C}_*(-)$ denotes the reduced singular chain functor, and " $\wedge$ " is the usual smash product of spaces, we have the following corollary.

Corollary 1.5. *For $d > 2\max\{m_i \mid 1 \leq i \leq r\} + 1$, there is a quasi-isomorphism*

$$C_*(\overline{\text{Emb}}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)) \otimes \mathbb{Q} \simeq \text{hRmod}_{\Omega}(\widetilde{C}_*((\bigvee_{i=1}^r S^{m_i})^{\wedge \bullet}, \mathbb{Q}), \widehat{H}_*(C(\bullet, \mathbb{R}^d), \mathbb{Q})).$$

To finish the proof of Theorem 0.1 we only need to show that the Ω -module $\widetilde{C}_*((\bigvee_{i=1}^r S^{m_i})^{\wedge \bullet})$ is rationally formal. This follows from Lemma 1.6 and also the fact that each $\widetilde{C}_*((S^{m_i})^{\wedge \bullet})$, $1 \leq i \leq r$, is formal (by [2, Lemma 8.5]).

Lemma 1.6. *Let X and Y be two pointed topological spaces. Assume that the right Ω -modules $\widetilde{C}_*(X^{\wedge \bullet})$ and $\widetilde{C}_*(Y^{\wedge \bullet})$ are formal in the category Rmod_{Ω} of right Ω -modules. Then $\widetilde{C}_*((X \vee Y)^{\wedge \bullet})$ is formal in the same category.*

Proof. Notice first that for each $k \geq 0$ we have

$$(X \vee Y)^{\wedge k} \simeq \bigvee_{f: \{1, \dots, k\} \rightarrow \{1, 2\}} X^{\wedge f^{-1}(1)} \wedge Y^{\wedge f^{-1}(2)}.$$

This implies

$$\begin{aligned} \widetilde{C}_*((X \vee Y)^{\wedge k}) &\simeq \bigoplus_{f: \{1, \dots, k\} \rightarrow \{1, 2\}} \widetilde{C}_*(X^{\wedge f^{-1}(1)} \wedge Y^{\wedge f^{-1}(2)}) \simeq \\ &\bigoplus_{f: \{1, \dots, k\} \rightarrow \{1, 2\}} \widetilde{C}_*(X^{\wedge f^{-1}(1)}) \otimes \widetilde{C}_*(Y^{\wedge f^{-1}(2)}). \end{aligned} \quad (1.6)$$

Consider the contravariant functor $\lambda: \Omega \times \Omega \longrightarrow \text{Ch}_*$, from $\Omega \times \Omega$ to chain complexes, defined by $\lambda(p, q) = \tilde{C}_*(X^{\wedge p}) \otimes \tilde{C}_*(Y^{\wedge q})$. Consider also the functor $\Delta: \text{Rmod}_{\Omega \times \Omega} \rightarrow \text{Rmod}_{\Omega}$ defined by

$$\Delta(\mu)(k) = \bigoplus_{f: \{1, \dots, k\} \rightarrow \{1, 2\}} \mu(f^{-1}(1), f^{-1}(2)),$$

where μ is a right $\Omega \times \Omega$ module. By (1.6), $\Delta(\lambda)$ is equivalent to $\tilde{C}_*((X \vee Y)^{\wedge \bullet})$ whose formality we want to show. Since λ is formal by hypothesis, that is there exists a zigzag of natural quasi-isomorphisms, $\lambda \simeq H_*(\lambda)$, between λ and $H_*(\lambda)$, it follows that

$$\Delta(\lambda) \simeq \Delta(H_*(\lambda)) = H_*(\Delta(\lambda)).$$

□

2 Hairy graph-complexes

The goal of this section is to prove Theorem 0.2 announced in the introduction.

2.1 Graph-complexes $\mathcal{E}_{\pi}^{m_1, \dots, m_r, d}$ and $\mathcal{E}_H^{m_1, \dots, m_r, d}$

In this section we will explicitly describe two graph-complexes: $\mathcal{E}_H^{m_1, \dots, m_r, d}$ and $\mathcal{E}_{\pi}^{m_1, \dots, m_r, d}$. In the next section we will show that they respectively compute the rational homology and the rational homotopy of $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$. Our description is exactly the same as that of complexes $\mathcal{E}_H^{m, n}$ and $\mathcal{E}_{\pi}^{m, n}$ introduced in [3, Section 2.1], except that our graphs have their external vertices "decorated" or "colored" with $\{1, \dots, r\}$ as the set of colours. As before the colours correspond to the components of the links.

Let us start with $\mathcal{E}_{\pi}^{m_1, \dots, m_r, d}$, which is a differential graded vector space spanned by some connected *hairy colored graphs* defined below, modulo certain relations. A *hairy graph* G is a triple (V_G^E, V_G^I, E_G) where

- V_G^E is a finite non-empty set whose elements are non-labeled *external* vertices of valence 1;
- V_G^I is a finite set (which can be empty) whose elements are non-labeled *internal* vertices of valence ≥ 3 ;
- E_G is the set of oriented edges, which may include tadpoles and multiple edges.

Notice that our graphs are connected. A *hairy colored graph* is a couple (G, f_G) where $f_G: V_G^E \longrightarrow \{1, \dots, r\}$ is a map from the set of external vertices to the set of colours. Given a colored graph G , an *orientation set* is the union $V_G^E \cup V_G^I \cup E_G$. Elements of V_G^E are assigned degree $-d$; elements of V_G^I of color i are assigned degree $-m_i$; elements of E_G are assigned degree $d - 1$. We will say that a colored graph is *oriented* if it is equipped with a total order O_G of its orientation set. Two graphs (G, f_G, O_G) and $(G', f_{G'}, O_{G'})$ are equivalent if there are two bijections

$$g: V_G^E \cup V_G^I = V_G \xrightarrow{\cong} V_{G'} = V_{G'}^E \cup V_{G'}^I \quad \text{and} \quad h: E_G \xrightarrow{\cong} E_{G'},$$

respecting adjacency, orientation of edges, order of orientation sets, and coloration. (Since g preserves the valence, it must send external vertices to external and internal to internal.) Let $\mathcal{G}$ be the collection of all equivalence classes of oriented colored graphs, and let $\mathcal{E}_{\pi}^{m_1, \dots, m_r, d}$ denote the vector space spanned by $\mathcal{G}$ modulo two relations in $\mathbb{Q}[\mathcal{G}]$:

- $(G, f_G, O_G) = (-1)^d(G', f_{G'}, O_{G'})$ if (G, f_G, O_G) differs from $(G', f_{G'}, O_{G'})$ only by an orientation of an edge;
- $(G, f_G, O_G) = \pm(G', f_{G'}, O_{G'})$ if $(G', f_{G'}, O_{G'})$ is obtained from (G, f_G, O_G) only by a permutation of the orientation set. The sign " $\pm$ " is the Koszul sign of permutation taking into account the degrees of elements.

Given a colored hairy graph (G, f_G) , we define its degree as a sum of degrees of the elements in its orientation set:

$$\text{degree}(G) = (d-1)|E_G| - d|V_G^I| - \sum_{i=1}^r m_i |f_G^{-1}(i)|.$$

This endows $\mathcal{E}_\pi^{m_1, \dots, m_r, d}$ with a grading. It is also equipped with a differential ∂ that turns it into a chain complex. The differential of a graph is defined as the sum of expansion of its internal vertices. More precisely, let (G, f_G, O_G) be an oriented colored graph. For any internal vertex $x \in V_G^I$ of valence l , let $(G_{px}, f_{G_{px}}, O_{G_{px}})$, $1 \leq p \leq n_x$, denote the graphs obtained from the expansion of x . One can see that the number of such graphs is $n_x = 2^{l-1} - l - 1$. They are defined as follows. Notice first that an expansion produces a new vertex and a new edge. For $1 \leq p \leq n_x$, one has

$$G_{px} = (V_{G_{px}}^E, V_{G_{px}}^I, E_{G_{px}}) = (V_G^E, V_G^I \cup \{v_{px}\}, E_G \cup \{e_{px}\}), \quad f_{G_{px}} = f_G,$$

and the orientation set has two more elements: v_{px} – the new vertex, and e_{px} – the new edge (oriented from the old vertex to the new one). Its order is defined as follows: $v_{px} \leq e_{px}$ and for all y in the orientation set of G , one has $e_{px} \leq y$ (here " $\leq$ " denotes the total order).

The differential is then defined by

$$\partial(G, f_G, O_G) = \sum_{x \in V_G^I} \sum_{p=1}^{n_x} (G_{px}, f_{G_{px}}, O_{G_{px}}).$$

Notice that there is no sign in each term of the differential. It is hidden in the way we order the orientation set and orient the new edge. Let us see one example. Assume that $r = 2$, that is, we have 2 colours, and consider the graph in Figure 1.




Figure 1

The three graphs that appear in the differential of Figure 1 are Figure 2, Figure 3 and Figure 4.

Now we define $\mathcal{E}_H^{m_1, \dots, m_r, d}$ as the cofree cocommutative coalgebra cogenerated by $\mathcal{E}_\pi^{m_1, \dots, m_r, d}$. We can view $\mathcal{E}_H^{m_1, \dots, m_r, d}$ as a vector space spanned by all possibly empty or disconnected colored hairy graphs (with each connected component from $\mathcal{E}_\pi^{m_1, \dots, m_r, d}$).




Figure 2




Figure 3




Figure 4

2.2 The rational homotopy of $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$

The goal of this section is to prove Theorem 0.2, which determines the rational homotopy of the space $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$. The plan of the proof is as follows. We will first prove that the graph-complex $\mathcal{E}_H^{m_1, \dots, m_r, d}$ is quasi-isomorphic to the complex $\text{hRmod}(\tilde{H}_*((\bigvee_{i=1}^r S^{m_i})^{\wedge \bullet}, \mathbb{Q}), \hat{H}_*(\mathcal{B}_d(\bullet), \mathbb{Q}))$ (where $\mathcal{B}_d$ denotes the little d -disks operad). The latter complex is the derived mapping space. Since both the source and target have trivial homology, the complex itself is equivalent to a product of Ext groups that can be either computed by taking the projective resolution of the source (as in Section 4) or by taking the injective resolution of the target. We show that the second approach produces exactly the complex $\mathcal{E}_H^{m_1, \dots, m_r, d}$. A similar argument shows that $\mathcal{E}_\pi^{m_1, \dots, m_r, d}$ is equivalent to the derived mapping space $\text{hRmod}_\Omega(\tilde{H}_*((\bigvee_{i=1}^r S^{m_i})^{\wedge \bullet}, \mathbb{Q}), \mathbb{Q} \otimes \hat{\pi}_* \mathcal{B}_d(\bullet))$. On the other hand, since $\mathcal{E}_H^{m_1, \dots, m_r, d}$ computes $H_*(\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \mathbb{Q})$, by dimensional reasons we conclude that $\mathcal{E}_\pi^{m_1, \dots, m_r, d}$ computes $\mathbb{Q} \otimes \pi_* \overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$. Which finishes the proof of Theorem 0.2.

In [3] Arone and Turchin explicitly describe a differential graded right Ω -modules $\hat{D}_d^\bullet$ that turns out to be an injective resolution of $\hat{H}_*(\mathcal{B}_d(\bullet), \mathbb{Q})$. We will first recall the right Γ -module $D_d^\bullet$, generated by certain graphs, introduced in [42]. For $k \geq 0$, D_d^k is a vector space spanned by possibly disconnected graphs with k external vertices labeled by $\{1, \dots, k\}$ of any valence ≥ 0 and some possibly empty set of non-labeled internal vertices of valence ≥ 3 , such that each their connected component has at least one external vertex. Tadpoles and multiple edges are allowed. The graphs are oriented, and an orientation set of such graph consists of the union of edges (considered as elements of degree $d-1$) and internal vertices (considered as elements of degree $-d$). Orienting a graph means putting a total order on its orientation set. As in Section 2.1, there is a similar notion of equivalence of graphs, and similar sign relations. Again as in Section 2.1, the differential is the sum of expansions of vertices (notice that, here, the expansion concerns both external and internal vertices). The expansion of an internal vertex is as in Section 2.1, while the expansion of an external vertex produces an external vertex (with the same label) and an internal vertex.

It is well known that the sequence $D_d^\bullet = \{D_d^k\}_{k \geq 0}$ is an operad (Figure 5 shows an example of the composition) in differential graded cocommutative coalgebras (Figure 6 is an example of the coproduct in D_d^3).

It is also well known [28, 42] that there is an inclusion of operads (in coalgebras) $H_*(\mathcal{B}_d(\bullet), \mathbb{Q}) \hookrightarrow D_d^\bullet$ that turns out to be a quasi-isomorphism. This inclusion implies that $D_d^\bullet$ is an infinitesimal bimodule over $H_*(\mathcal{B}_d(\bullet), \mathbb{Q})$, and therefore over $H_0(\mathcal{B}_d(\bullet), \mathbb{Q}) = \text{Com}$. By [2, Proposition 4.9], we deduce that $D_d^\bullet$ is a right Γ -module. Moreover, since the right Γ -module structure respects the coalgebra structure, $D_d^\bullet$ is a right Γ -module in coalgebras. Recalling the cross effect functor cr from (1.5), one defines the right Ω -module $\hat{D}_d^\bullet$




Figure 5: Examples of the insertion operation




Figure 6: An example of the coproduct in D_d^3

by

$$\hat{D}_d^\bullet = cr(D_d^\bullet).$$

It is easy to see (by the definition of the cross effect) that for each $k \geq 0$, $\hat{D}_d^k$ is spanned by the same generators as D_d^k , except that generators of $\hat{D}_d^k$ are graphs whose external vertices are of valence ≥ 1 . The following result is [3, Proposition 2.6].

Proposition 2.1. *The right Ω -module $\hat{D}_d^\bullet$ is finite dimensional and injective in each homological degree. The inclusion $\hat{H}_*(\mathcal{B}_d(\bullet), \mathbb{Q}) \hookrightarrow \hat{D}_d^\bullet$ is a quasi-isomorphism of Ω -modules and thus is an injective resolution for every given complexity.*

Remark 2.2. *To recall the decomposition by complexity t in $\hat{H}_*(\mathcal{B}_d(\bullet), \mathbb{Q})$ is $\bigoplus_t \hat{H}_{t(d-1)}(\mathcal{B}_d(\bullet), \mathbb{Q})$. In $\hat{D}_d^\bullet$ one defines a similar splitting saying that a complexity of a graph $G \in \hat{D}_d^k$ is the first Betti number of the graph obtained from G by gluing together all external vertices.*

Theorem 2.3. *For $d > 2\max\{m_i \mid 1 \leq i \leq r\} + 1$ the graph-complex $\mathcal{E}_H^{m_1, \dots, m_r, d}$ is quasi-isomorphic to $\text{hRmod}_\Omega(\tilde{H}_*((\bigvee_{i=1}^r S^{m_i})^{\wedge \bullet}, \mathbb{Q}), \hat{H}_*(\mathcal{B}_d(\bullet), \mathbb{Q}))$, that is,*

$$\mathcal{E}_H^{m_1, \dots, m_r, d} \simeq \text{hRmod}_\Omega(\tilde{H}_*((\bigvee_{i=1}^r S^{m_i})^{\wedge \bullet}, \mathbb{Q}), \hat{H}_*(\mathcal{B}_d(\bullet), \mathbb{Q})). \quad (2.1)$$

Proof. The steps of the proof are the same as those of the proof of Theorem 2.2 from [3]. Thus, for certain details, we refer the reader to [3]. For each $k \geq 0$, let $M(D_d^k)$ denote the subspace of D_d^k generated by graphs

whose external vertices are of valence = 1. We have the following quasi-isomorphisms

$$\begin{aligned}
& \text{hRmod}_{\Omega}(\tilde{H}_*((\bigvee_{i=1}^r S^{m_i})^{\wedge \bullet}, \mathbb{Q}), \hat{H}_*(\mathcal{B}_d(\bullet), \mathbb{Q})) \\
& \simeq \text{Rmod}_{\Omega}(\tilde{H}_*((\bigvee_{i=1}^r S^{m_i})^{\wedge \bullet}, \mathbb{Q}), \hat{D}_d^{\bullet}) \text{ by Proposition 2.1} \\
& \simeq \bigoplus_{k=0}^{\infty} \text{hom}_{\Sigma_k} \left(\tilde{H}_*((\bigvee_{i=1}^r S^{m_i})^{\wedge k}, \mathbb{Q}), M(D_d^k) \right) \text{ by [3, Lemma 2.7 and Lemma 2.8]} \\
& \simeq \bigoplus_{k=0}^{\infty} \text{hom}_{\Sigma_k} \left(\bigoplus_{s_1+\dots+s_r=k} \text{Ind}_{\Sigma_{s_1} \times \dots \times \Sigma_{s_r}}^{\Sigma_k} \tilde{H}_*(S^{s_1 m_1 + \dots + s_r m_r}, \mathbb{Q}), M(D_d^k) \right) \text{ by (0.3)} \\
& \simeq \bigoplus_{k=0}^{\infty} \bigoplus_{s_1+\dots+s_r=k} \text{hom}_{\Sigma_{s_1} \times \dots \times \Sigma_{s_r}} \left(\tilde{H}_*(S^{s_1 m_1 + \dots + s_r m_r}, \mathbb{Q}), M(D_d^k) \right).
\end{aligned}$$

By the description of the graph-complex $\mathcal{E}_H^{m_1, \dots, m_r, d}$ from Section 2.1, it is not difficult to see that there is an isomorphism of chain complexes

$$\mathcal{E}_H^{m_1, \dots, m_r, d} \cong \bigoplus_{k=0}^{\infty} \bigoplus_{s_1+\dots+s_r=k} \text{hom}_{\Sigma_{s_1} \times \dots \times \Sigma_{s_r}} \left(\tilde{H}_*(S^{s_1 m_1 + \dots + s_r m_r}, \mathbb{Q}), M(D_d^k) \right),$$

which completes the proof. $\square$

One should mention that the decomposition over $(s_1, \dots, s_r)$ above appeared exactly as the multidegree Hodge splitting.

Remark 2.4. *A careful reader might prefer to see a product instead of the direct sum used in the above equations. But it is easy to show that $\mathcal{E}_H^{m_1, \dots, m_r, d}$ is finite dimensional in every homological degree, for which reason it would not matter whether we take a product or a direct sum of its components, see [2, Remark 13.3] and [3, Section 5] for degree estimates in similar graph-complexes.*

The following corollary is an immediate consequence of Theorem 0.1 and Theorem 2.3.

Corollary 2.5. *For $d > 2\max\{m_i \mid 1 \leq i \leq r\} + 1$ the graph-complex $\mathcal{E}_H^{m_1, \dots, m_r, d}$ computes the rational homology of the space $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$, that is,*

$$H(\mathcal{E}_H^{m_1, \dots, m_r, d}) \cong H_*(\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \mathbb{Q}). \quad (2.2)$$

Corollary 2.6. *For $d > 2\max\{m_i \mid 1 \leq i \leq r\} + 1$ the graph-complex $\mathcal{E}_{\pi}^{m_1, \dots, m_r, d}$ computes the rational homotopy of the space $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$, that is,*

$$H(\mathcal{E}_{\pi}^{m_1, \dots, m_r, d}) \cong \mathbb{Q} \otimes \pi_*(\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)). \quad (2.3)$$

Proof. The desired result is a combination of Corollary 2.5 and the following two facts:

- the rational homology of the space $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$ is a graded polynomial coalgebra cogenerated by the rational homotopy of the same space;

- the subcomplex of $\widehat{\mathcal{E}}_H^{m_1, \dots, m_r, d}$, formed by primitive elements, is exactly the complex $\mathcal{E}_\pi^{m_1, \dots, m_r, d}$.

Note that these two facts imply that (2.2) and (2.3) are equivalent. $\square$

Theorem 2.7. *For $d > 2\max\{m_i \mid 1 \leq i \leq r\} + 1$ the graph-complex $\mathcal{E}_\pi^{m_1, \dots, m_r, d}$ is quasi-isomorphic to $\mathrm{hRmod}(\widetilde{H}_*((\bigvee_{i=1}^r S^{m_i})^{\wedge \bullet}, \mathbb{Q}), \mathbb{Q} \otimes \widehat{\pi}_* \mathcal{B}_d(\bullet))$.*

Proof. The proof goes in the same way as that of Theorem 2.3. Instead of $\widehat{D}_d^\bullet$, one constructs a complex of injective Ω -modules $\widehat{P}_d^\bullet \simeq \mathbb{Q} \otimes \widehat{\pi}_* \mathcal{B}_d(\bullet)$ (see [3, Proposition 2.4] or [42]) defined as the cross effect of the right Γ -module $P_d^\bullet = \{P_d^k\}_{k \geq 0}$. Here P_d^k is the primitive part of D_d^k – it is spanned by so called *internally connected graphs*, that have exactly one connected component different from singleton. It is also important that the quasi-isomorphism $\widehat{P}_d^\bullet \simeq \mathbb{Q} \otimes \widehat{\pi}_* \mathcal{B}_d(\bullet)$ respects the splitting by complexity t . In $\widehat{P}_d^\bullet \subset \widehat{D}_d^\bullet$ it is recalled in Remark 2.2, while for the right-hand side it is $\mathbb{Q} \otimes \widehat{\pi}_* \mathcal{B}_d(\bullet) \simeq \bigoplus_t \mathbb{Q} \otimes \widehat{\pi}_{t(d-2)+1} \mathcal{B}_d(\bullet)$. $\square$

We are now ready to prove Theorem 0.2.

Proof of Theorem 0.2. It is an immediate consequence of Corollary 2.6 and Theorem 2.7. $\square$

3 Low codimensional high dimensional string links

This section is devoted to Conjecture 3.1. We explain here what supports this conjecture and why we think it is true.

Conjecture 3.1. *The statement (2.3) of Corollary 2.6 holds for any codimension $d - \max\{m_i \mid 1 \leq i \leq r\} > 2$, and any non-negative degree $*$ ≥ 0 .*

There are two things that support this conjecture. Firstly, it is true for $r = 1$, which was recently shown by Fresse, Willwacher, and Turchin [16]. Secondly, Theorem 3.2 below shows that it is always true on the level of π_0 .

As for the case $r = 1$, we expect two main ingredients in a possible proof of the conjecture: the relative Hopf formality of the little discs operads, which luckily holds in any codimension ≥ 2 [44], and certain delooping statement for $\overline{\mathrm{Emb}}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$, which would say that $\overline{\mathrm{Emb}}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$ is an iterated loop space of the derived mapping space between two algebraic objects described in terms of the little discs operads. Unfortunately, the problem of delooping in the case $r \geq 2$ appears to be much more difficult. An attempt for such delooping in the classical case of string links (all $m_i = 1$) has been done in [13]. It describes an interesting possible candidate of the delooping, but unfortunately, the paper has a mistake in the proof of its main result.

Theorem 3.2. *For $d - \max\{m_i \mid 1 \leq i \leq r\} > 2$, $\pi_0 \overline{\mathrm{Emb}}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$, with the product induced by the concatenation of links, is a finitely generated abelian group. Moreover, one has*

$$\mathbb{Q} \otimes \pi_0 \overline{\mathrm{Emb}}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) \simeq H_0 \mathcal{E}_\pi^{m_1, \dots, m_r, d}. \quad (3.1)$$

One should mention that the fact that $\pi_i \overline{\text{Emb}}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$, $i > 0$, are finitely generated abelian groups, easily follows from the embedding calculus, which is always applicable when codimension > 2 [19, 20].

This theorem is a consequence of Lemmas 3.3 and 3.8. Let us first understand the right-hand side of (3.1). The complex $\mathcal{E}_\pi^{m_1, \dots, m_r, d}$ splits into a direct sum of complexes

$$\mathcal{E}_\pi^{m_1, \dots, m_r, d} = \bigoplus_{g \geq 0} \mathcal{E}_{g\pi}^{m_1, \dots, m_r, d},$$

where $\mathcal{E}_{g\pi}^{m_1, \dots, m_r, d}$ is spanned by the graphs whose first Betti number (= genus) is g .

Lemma 3.3. *For $d - \max\{m_i \mid 1 \leq i \leq r\} > 2$, $H_{\leq 0} \mathcal{E}_\pi^{m_1, \dots, m_r, d} = H_{\leq 0} \mathcal{E}_{0\pi}^{m_1, \dots, m_r, d}$.*

Proof. It is enough to show that all the graphs of positive genus have positive homological degree. One can proceed by induction. The degree of any graph of the form  is $d - m_i - 2 > 0$. Any uni-trivalent hairy graph is obtained from  by a sequence of attachments of hairs (inside an edge) and edges (both endpoints on the same edge or on two different ones). The first operation creates 2 new edges and two vertices – one external and one internal. The total degree change is

$$2(d-1) - d - m_i = d - m_i - 2 > 0.$$

The second operation creates 3 new edges and two new internal vertices. The total degree change is

$$3(d-1) - 2d = d - 3 > 0.$$

Finally, any other hairy graph is obtained from a uni-trivalent one by a sequence of contractions of edges. Each such contraction increases the degree by one. $\square$

In case $r = 1$, $H_* \mathcal{E}_{0\pi}^{m_1, d}$ is always one-dimensional. In case of even codimension $d - m_1$, it is spanned by the line graph  of degree $d - 2m_1 - 1$; in case of odd codimension – by the tripod  of degree $2d - 2m_1 - 3$. The following result was obtained in [16] as a corollary of (3.1), which was proved there for $r = 1$.

Corollary 3.4 ([16]). *For $d - m > 2$, $\pi_0 \overline{\text{Emb}}_c(\mathbb{R}^m, \mathbb{R}^d)$ is a finitely generated abelian group of rank ≤ 1 . It is infinite only if either $m = 2k + 1$, $d = 4k + 3$, $k \geq 1$, or $m = 4k - 1$, $d = 6k$, $k \geq 1$.*

One has a fiber sequence

$$\overline{\text{Emb}}_c(\mathbb{R}^m, \mathbb{R}^d) \rightarrow \text{Emb}_c(\mathbb{R}^m, \mathbb{R}^d) \rightarrow \Omega^m V_m(\mathbb{R}^d), \quad (3.2)$$

where $V_m(\mathbb{R}^d)$ is the Stiefel manifold $SO(d)/SO(d-m)$. (By the Smale-Hirsch principle $\text{Imm}_c(\mathbb{R}^m, \mathbb{R}^d) \simeq \Omega^m V_m(\mathbb{R}^d)$). From (3.2) we get the long exact sequence

$$\dots \rightarrow \pi_{m+1} V_m(\mathbb{R}^d) \xrightarrow{j_*} \pi_0 \overline{\text{Emb}}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) \xrightarrow{k_*} \pi_0 \text{Emb}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) \xrightarrow{l_*} \pi_m V_m(\mathbb{R}^d). \quad (3.3)$$

The case $m = 2k + 1$, $d = 4k + 3$, $k \geq 1$, of the corollary corresponds to the graph-cycle . Geometrically it appears as the image under j_* of the Euler class of $SO(d-m)$ in $V_m(\mathbb{R}^d) = SO(d)/SO(d-m)$. The case $m = 4k - 1$, $d = 6k$, $k \geq 1$, corresponds to the graph-cycle . Its image in $\pi_0 \text{Emb}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$ under

k_* is the Haefliger trefoil [23], which is known to be trivial as immersion (equivalently lies in the image of k_*).

To describe $H_0\mathcal{E}_{0\pi}^{m_1, \dots, m_r, d}$ we recall some standard facts and notation.

The operad Lie is cyclic, thus $\text{Lie}(n-1)$ is a Σ_n -module. The following result is well known.

Lemma 3.5 ([29, 22, 9]). *For $n \geq 1$, one has*

$$\text{Lie}(n-1) \simeq_{\Sigma_n} \ker \left(\text{Ind}_{\Sigma_{n-1}}^{\Sigma_n} \text{Lie}(n-1) \xrightarrow{q} \text{Lie}(n) \right),$$

where the map q is described below.

Elements of $\text{Ind}_{\Sigma_{n-1}}^{\Sigma_n} \text{Lie}(n-1)$ can be written in the form $\sum_{i=1}^n \alpha_i x_i \otimes f_i(x_1, \dots, \hat{x}_i, \dots, x_n)$, where $\alpha_i \in \mathbb{Q}$, $f_i \in \text{Lie}(n-1)$. The map q sends

$$q: x_i \otimes f_i(x_1, \dots, \hat{x}_i, \dots, x_n) \mapsto [x_i, f_i(x_1, \dots, \hat{x}_i, \dots, x_n)].$$

For any symmetric sequence $M(\bullet)$ and any graded vector space V , we denote by

$$F_{M(\bullet)}(V) := \bigoplus_{k \geq 0} M(k) \otimes_{\Sigma_k} V^{\otimes k}.$$

Lemma 3.6. *For any $d - \max\{m_i \mid 1 \leq i \leq r\} > 2$,*

$$H_*\mathcal{E}_{0\pi}^{m_1, \dots, m_r, d} = \Sigma^{-d+3} F_{\text{Lie}(\bullet-1)}(\tilde{H}_*(\bigvee_{i=1}^r S^{d-m_i-2}, \mathbb{Q})).$$

Proof. It is well known that the homology of the complex of trees, whose leaves are bijectively labeled by $\{1, \dots, n\}$, is concentrated in the top homological degree and (up to a regrading and a possible tensor with the sign representation) is isomorphic to $\text{Lie}(n-1)$, see [47]. A careful count of the degrees and signs finishes the proof. $\square$

In particular, this lemma says that $H_0\mathcal{E}_{0\pi}^{m_1, \dots, m_r, d}$ is isomorphic to the degree $d-3$ part of $F_{\text{Lie}(\bullet-1)}(\tilde{H}_*(\bigvee_{i=1}^r S^{d-m_i-2}, \mathbb{Q}))$.

Now we want to concentrate on the left-hand side of (3.1). Let $\text{Emb}(\coprod_{i=1}^r S^{m_i}, S^d)$ be the space of smooth spherical links $\coprod_{i=1}^r S^{m_i} \hookrightarrow S^d$. It was shown by A. Haefliger [24] that for $d - \max\{m_i \mid 1 \leq i \leq r\} > 2$, $\pi_0 \text{Emb}(\coprod_{i=1}^r S^{m_i}, S^d)$ is a finitely generated abelian group, whose product is defined as follows. Given two links $L_1, L_2 \in \text{Emb}(\coprod_{i=1}^r S^{m_i}, S^d)$, we move one inside one semisphere, and the other inside the complement semisphere. Then we connect the components of L_1 with the corresponding components of L_2 by very narrow tubes to form a new link. The result as an element of $\pi_0 \text{Emb}(\coprod_{i=1}^r S^{m_i}, S^d)$ does not depend on the paths of the tubes since the complement of $L_1 \amalg L_2$ is simply connected. By $\pi_0^U \text{Emb}(\coprod_{i=1}^r S^{m_i}, S^d)$, $\pi_0^U \text{Emb}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$, $\pi_0^U \overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$ we denote the kernel of the corresponding map

$$\begin{aligned} \pi_0 \text{Emb}(\coprod_{i=1}^r S^{m_i}, S^d) &\rightarrow \prod_{i=1}^r \pi_0 \text{Emb}(S^{m_i}, S^d); \\ \pi_0 \text{Emb}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) &\rightarrow \prod_{i=1}^r \pi_0 \text{Emb}_c(\mathbb{R}^{m_i}, \mathbb{R}^d); \\ \pi_0 \overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) &\rightarrow \prod_{i=1}^r \pi_0 \overline{\text{Emb}}_c(\mathbb{R}^{m_i}, \mathbb{R}^d). \end{aligned}$$

The letter U refers to “unknot”. Each of the groups is the subgroup of links formed by unknots.

The following lemma proves the first statement of Theorem 3.2. Its proof is an easy exercise, which we leave to the reader.

Lemma 3.7. *For $d - \max\{m_i \mid 1 \leq i \leq r\} > 2$, one has isomorphisms of monoids (and thus of finitely generated abelian groups):*

- $\pi_0 \text{Emb}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) \simeq \pi_0 \text{Emb}(\coprod_{i=1}^r S^{m_i}, S^d);$
- $\pi_0^U \overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) \simeq \pi_0^U \text{Emb}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) \simeq \pi_0^U \text{Emb}(\coprod_{i=1}^r S^{m_i}, S^d);$
- $\pi_0 \overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) \simeq \pi_0^U \overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) \times \prod_{i=1}^r \pi_0 \overline{\text{Emb}}_c(\mathbb{R}^{m_i}, \mathbb{R}^d).$

By [24] and Corollary 3.4, $\pi_0 \text{Emb}(\coprod_{i=1}^r S^{m_i}, S^d)$ and $\pi_0 \overline{\text{Emb}}_c(\mathbb{R}^{m_i}, \mathbb{R}^d)$ are finitely generated abelian groups, thus $\pi_0 \overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$ is so as well.

Define the groups

$$\begin{aligned}\overline{\Pi}^{m_1 \dots m_r d} &:= \pi_{d-2}(\vee_{i=1}^r S^{d-m_i-1}); \\ \overline{\Lambda}^{m_1 \dots m_r d} &:= \bigoplus_{i=1}^r \pi_{m_i}(\vee_{i=1}^r S^{d-m_i-1}); \\ \overline{K}^{m_1 \dots m_r d} &:= \ker(\overline{\Lambda}^{m_1 \dots m_r d} \xrightarrow{q_0} \overline{\Pi}^{m_1 \dots m_r d}),\end{aligned}$$

here $q_0: (\alpha_1 \dots \alpha_r) \mapsto \sum_{i=1}^r [\iota_i, \alpha_i]$, where $[\cdot, \cdot]$ is the Whitehead bracket, $\iota_i \in \pi_{d-m_i-1}(\vee_{i=1}^r S^{d-m_i-1})$ is the element encoding the inclusion of the i -th summand $S^{d-m_i-1} \hookrightarrow \vee_{i=1}^r S^{d-m_i-1}$. We also define their subgroups

$$\begin{aligned}\Pi^{m_1 \dots m_r d} &:= \ker \left(\pi_{d-2}(\vee_{i=1}^r S^{d-m_i-1}) \rightarrow \bigoplus_{i=1}^r \pi_{d-2} S^{d-m_i-1} \right); \\ \Lambda^{m_1 \dots m_r d} &:= \bigoplus_{i=1}^r \ker \left(\pi_{m_i}(\vee_{i=1}^r S^{d-m_i-1}) \rightarrow \pi_{m_i} S^{d-m_i-1} \right); \\ K^{m_1 \dots m_r d} &:= \ker(\Lambda^{m_1 \dots m_r d} \xrightarrow{q_0|_{\Lambda}} \Pi^{m_1 \dots m_r d}).\end{aligned}$$

Lemma 3.8. *For $d - \max\{m_i \mid 1 \leq i \leq r\} > 2$,*

$$\mathbb{Q} \otimes \pi_0 \overline{\text{Emb}}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) \simeq \mathbb{Q} \otimes \overline{K}^{m_1 \dots m_r d} \simeq H_0 \mathcal{E}_{0\pi}^{m_1, \dots, m_r, d}. \quad (3.4)$$

Proof. We prove below the second isomorphism in (3.4) for any $r \geq 1$. Thus, by Corollary 3.4 it's enough to show the first isomorphism only for $r \geq 2$. In [24], A. Haefliger develops a homotopical approach to compute $\pi_0 \text{Emb}(\coprod_{i=1}^r S^{m_i}, S^d)$. Using this method it was shown in [10, Lemma 1.3] that

$$\mathbb{Q} \otimes \pi_0^U \text{Emb}(\prod_{i=1}^r S^{m_i}, S^d) = \mathbb{Q} \otimes K^{m_1 \dots m_r d}.$$

Applying Lemma 3.7, we get

$$\begin{aligned} \mathbb{Q} \otimes \pi_0 \overline{\text{Emb}}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) &\simeq \mathbb{Q} \otimes \pi_0^U \overline{\text{Emb}}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) \oplus \bigoplus_{i=1}^r \mathbb{Q} \otimes \pi_0 \overline{\text{Emb}}_c(\mathbb{R}^{m_i}, \mathbb{R}^d) \\ &\simeq \mathbb{Q} \otimes K^{m_1 \dots m_r d} \oplus \bigoplus_{i=1}^r \mathbb{Q} \otimes \overline{K}^{m_i d} \simeq \mathbb{Q} \otimes \overline{K}^{m_1 \dots m_r d}. \end{aligned}$$

The last isomorphism is due to the splittings

$$\begin{aligned} \overline{\Pi}^{m_1 \dots m_r d} &= \Pi^{m_1 \dots m_r d} \oplus \bigoplus_{i=1}^r \pi_{d-2} S^{d-m_i-1} = \Pi^{m_1 \dots m_r d} \oplus \bigoplus_{i=1}^r \overline{\Pi}^{m_i d}; \\ \overline{\Lambda}^{m_1 \dots m_r d} &= \Lambda^{m_1 \dots m_r d} \oplus \bigoplus_{i=1}^r \pi_{m_i} S^{d-m_i-1} = \Lambda^{m_1 \dots m_r d} \oplus \bigoplus_{i=1}^r \overline{\Lambda}^{m_i d}, \end{aligned}$$

and also the fact that q_0 sends each summand of $\overline{\Lambda}^{m_1 \dots m_r d}$ to the corresponding summand of $\overline{\Pi}^{m_1 \dots m_r d}$.

To see the second isomorphism of (3.4), we notice that

$$\mathbb{Q} \otimes \pi_*(\bigvee_{i=1}^r S^{d-m_i-1}) \simeq \Sigma F_{\text{Lie}(\bullet)}(\tilde{H}_*(\bigvee_{i=1}^r S^{d-m_i-2}, \mathbb{Q})).$$

Thus $\mathbb{Q} \otimes \overline{\Pi}^{m_1 \dots m_r d}$ is the degree $d-3$ part of $F_{\text{Lie}(\bullet)}(\tilde{H}_*(\bigvee_{i=1}^r S^{d-m_i-2}, \mathbb{Q}))$. Similarly, one can see that $\mathbb{Q} \otimes \overline{\Lambda}^{m_1 \dots m_r d}$ is the degree $d-3$ part of $F_{\text{Ind}_{\Sigma_{\bullet-1}}^{\Sigma_{\bullet}} \text{Lie}(\bullet-1)}(\tilde{H}_*(\bigvee_{i=1}^r S^{d-m_i-2}, \mathbb{Q}))$. Thus $\overline{K}^{m_1 \dots m_r d}$ is the degree $d-3$ part of

$$\ker \left(F_{\text{Ind}_{\Sigma_{\bullet-1}}^{\Sigma_{\bullet}} \text{Lie}(\bullet-1)}(\tilde{H}_*(\bigvee_{i=1}^r S^{d-m_i-2}, \mathbb{Q})) \xrightarrow{F_q} F_{\text{Lie}(\bullet)}(\tilde{H}_*(\bigvee_{i=1}^r S^{d-m_i-2}, \mathbb{Q})) \right),$$

which by Lemmas 3.5 and 3.6 is the same as $H_0 \mathcal{E}_{0\pi}^{m_1, \dots, m_r, d}$.

□

4 The Koszul complexes

By Theorems 0.1 and 0.2, the rational homology and homotopy of $\overline{\text{Emb}}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$ is described in terms of derived maps of right Ω -modules. In this section we exhibit explicit complexes computing the (co)homology and homotopy of $\overline{\text{Emb}}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$ by considering the projective resolution of the source Ω -modules in (0.8) and (0.8). We call these complexes *Koszul* as they are constructed using operadic language and the Koszul duality between the commutative and Lie operads. At the end of the section we show that the Koszul complexes can also be used to understand rationally the stages of the Goodwillie-Weiss towers for $\overline{\text{Emb}}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$.

For $s_1, \dots, s_r \geq 0$, recall the definition of the right Ω -module $Q_{s_1 \dots s_r}^{m_1 \dots m_r}$ (it was introduced in (0.5)).

$$Q_{s_1 \dots s_r}^{m_1 \dots m_r}(k) = \begin{cases} 0 & \text{if } k \neq s_1 + \dots + s_r \\ \text{Ind}_{\Sigma_{s_1} \times \dots \times \Sigma_{s_r}}^{\Sigma_k} \tilde{H}_*(S^{s_1 m_1 + \dots + s_r m_r}; \mathbb{Q}) & \text{if } k = s_1 + \dots + s_r \end{cases} \quad (4.1)$$

Proposition 4.1. *For $d > 2\max\{m_i \mid 1 \leq i \leq r\} + 1$, one has quasi-isomorphisms*

$$C_*(\overline{\text{Emb}}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \otimes \mathbb{Q}) \simeq \bigoplus_{s_1, \dots, s_r, t \geq 0} \text{hRmod}_{\Omega}(Q_{s_1 \dots s_r}^{m_1 \dots m_r}, \widehat{H}_{t(d-1)}(C(\bullet, \mathbb{R}^d); \mathbb{Q})) \quad (4.2)$$

$$\pi_* \overline{\text{Emb}}_c(\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) \otimes \mathbb{Q} \simeq \bigoplus_{s_1, \dots, s_r, t \geq 0} \text{hRmod}_{\Omega}(Q_{s_1 \dots s_r}^{m_1 \dots m_r}, \widehat{\pi}_{t(d-2)+1} C(\bullet, \mathbb{R}^d) \otimes \mathbb{Q}) \quad (4.3)$$

Proof. This is a direct consequence of Theorems 0.1 and 0.2 and the fact that both the source and target Ω -modules in (0.1) and 0.2 split into a direct sum/product. The only thing we need to understand is why the product can be replaced by the direct sum. This follows for example, from Remark 2.4. $\square$

4.1 The Koszul dual of the right Ω -module $Q_{s_1 \dots s_r}^{m_1 \dots m_r}$

In this section we will find a cofibrant replacement (the model structure we use in the category Rmod_{Ω} is the one whose weak equivalences and fibrations are levelwise), denoted by $CQ_{s_1 \dots s_r}^{m_1 \dots m_r}$, of $Q_{s_1 \dots s_r}^{m_1 \dots m_r}$. We will follow exactly the same steps as in [3, Section 5 and Section 6], where this was done for $r = 1$. This cofibrant model will furnish us the so called *Koszul dual* of $Q_{s_1 \dots s_r}^{m_1 \dots m_r}$, which will be denoted by $KQ_{s_1 \dots s_r}^{m_1 \dots m_r}$. We will also explicitly describe $CQ_{s_1 \dots s_r}^{m_1 \dots m_r}$ (and then explicitly describe $KQ_{s_1 \dots s_r}^{m_1 \dots m_r}$) in terms of certain spaces of colored forests.

Let us start with an explicit description of $CQ_{s_1 \dots s_r}^{m_1 \dots m_r}$. Notice first that when $r = 1$, $Q_{s_1}^{m_1}$ is exactly the right Ω -module Q_s^m that appears in [3, Section 5]. Notice also that CQ_s^m is explicitly described in that section in terms of forests. Here we will describe $CQ_{s_1 \dots s_r}^{m_1 \dots m_r}$ in a similar way, except that in our case we will get "colored forests" with $\{1, \dots, r\}$ as the set of colours. The main idea of this construction is that right Ω -module is the same thing as the right module over the operad Com_+ of commutative non-unital algebras (that is, Com_+ is the commutative operad with zero in arity zero).

Recall first two standard notations. The first one is the operation " $\circ$ " (explicitly defined in [30, Chapter 5]) in the category of symmetric sequences in chain complexes [30, § 5.1]. Intuitively, if $M = \{M(n)\}_{n \geq 0}$ and $N = \{N(n)\}_{n \geq 0}$ are two symmetric sequences, elements in $M \circ N(n)$ are of the form $x(y_1, \dots, y_k)$, where $x \in M(k)$, $y_i \in N(i_k)$, and $i_1 + \dots + i_k = n$ (one should think about an element in $M(k)$ as an operation of arity k). Note that one can define " $\circ$ " in the category of symmetric sequences in any cocomplete symmetric monoidal category. This operation defines a monoidal structure on symmetric sequences. The unit for this structure is the sequence $\mathbf{1}$, defined as $\mathbf{1}(1)$ is the unit and every $\mathbf{1}(n)$, $n \neq 1$, is the initial element. Notice also that monoids with respect to $\circ$ are operads. The second notation is that of suspension of a symmetric sequence. Let $M = \{M(n)\}_{n \geq 0}$ be a symmetric sequence in chain complexes. The *operadic suspension*, denoted by $M[1]$, of M is a symmetric sequence defined by

$$M[1](n) = \Sigma^{n-1} M(n) \otimes \text{sign}_n,$$

where Σ^{n-1} is the usual $(n-1)$ -suspension, and sign_n is the sign representation of Σ_n . Notice that an operadic suspension of an operad is again an operad because of the identities $\mathbf{1}[1] = \mathbf{1}$ and $(M \circ N)[1] = M[1] \circ N[1]$.

Given a right module $\mathcal{R}$ over any Koszul operad $\mathcal{K}$, Fresse defines in [15] a cofibrant replacement, denoted by $C\mathcal{R}$, of $\mathcal{R}$. If $\mathcal{K}^!$ denotes the cooperad Koszul dual to $\mathcal{K}$, as a symmetric sequence $C\mathcal{R} = \mathcal{R} \circ \mathcal{K}^![1] \circ \mathcal{K}$. Its

differential consists of three summands. The first one is induced by the internal differential on $\mathcal{R}$. The second one arises from the right $\mathcal{K}$ module structure of $\mathcal{R}$. The third one arises from the left $\mathcal{K}$ module structure of $\mathcal{K}$. Notice that $C\mathcal{R}$ as a right $\mathcal{K}$ module is freely generated by the symmetric sequence $K\mathcal{R} = \mathcal{R} \circ \mathcal{K}$. This sequence $K\mathcal{R}$ with its natural differential is sometimes called the Koszul dual of $\mathcal{R}$. Notice that it naturally carries the structure of a right $\mathcal{K}^i[1]$ comodule. Applying Fresse's construction to $Q_{s_1 \dots s_r}^{m_1 \dots m_r}$, viewed as a right module over Com_+ (here the action of Com_+ on $Q_{s_1 \dots s_r}^{m_1 \dots m_r}$ is trivial), and by noticing that the cooperad Koszul dual of Com_+ is coLie , we get

$$CQ_{s_1 \dots s_r}^{m_1 \dots m_r} = Q_{s_1 \dots s_r}^{m_1 \dots m_r} \circ \text{coLie}[1] \circ \text{Com}_+, \quad (4.4)$$


where coLie is the dual of the operad $\mathcal{Lie}$ of Lie algebras. For obvious reasons only the third summand in the differential will be non-trivial. The equation (4.4) allows us to explicitly describe the right Ω -module $CQ_{s_1 \dots s_r}^{m_1 \dots m_r}$. Recall that the cooperad coLie can be described as certain space of trees modulo Arnold (3 term) relations [39]. Let A be a finite set of cardinal k . Then $CQ_{s_1 \dots s_r}^{m_1 \dots m_r}(A)$ or simply $CQ_{s_1 \dots s_r}^{m_1 \dots m_r}(k)$ is a chain complex spanned by the following set $\mathcal{F}_{s_1 \dots s_r}^{m_1 \dots m_r}(k)$ of *oriented colored forests*:

- A *vertex* in such forest F is a nonempty subset of A . The set of vertices forms a partition $\coprod_i A_i$ of A .
- Vertices of F , labeled by subsets A_i , are connected by edges so that no cycles appear, or, in other words, the obtained graph is a forest.
- Such forest F must consist of exactly $s_1 + \dots + s_r$ trees. We denote by T_F the set of its connected components.
- A *colored forest* is a couple (F, f) , where F is a forest as above, and $f: T_F \rightarrow \{1, \dots, r\}$ is a map (called coloring) from the set of trees in F to the set of colours $\{1, \dots, r\}$ such that for all $i = 1, \dots, r$, $|f^{-1}(i)| = s_i$.
- An *oriented colored forest* or an element in $\mathcal{F}_{s_1 \dots s_r}^{m_1 \dots m_r}(k)$ is a triple (F, f, O) , where (F, f) is a colored forest, and $O = T_F \cup E_F$ (here E_F denotes the set of all edges in F) is endowed with a total order $\leq$. The set O is the so called *orientation set*.

We define a degree of an element of $\mathcal{F}_{s_1 \dots s_r}^{m_1 \dots m_r}(k)$. Let (F, f, O) be any element in $\mathcal{F}_{s_1 \dots s_r}^{m_1 \dots m_r}(k)$. Each edge of F is of degree 1. A tree T in F is of degree m_i if $T \in f^{-1}(i)$. The total degree of (F, f, O) is then

$$\deg(F, f, O) = |E_F| + \sum_{1 \leq i \leq r} s_i m_i. \quad (4.5)$$

Let $\mathbb{Q}[\mathcal{F}_{s_1 \dots s_r}^{m_1 \dots m_r}(k)]$ denote the graded vector space over $\mathbb{Q}$ spanned by $\mathcal{F}_{s_1 \dots s_r}^{m_1 \dots m_r}(k)$. We consider two relations in $\mathbb{Q}[\mathcal{F}_{s_1 \dots s_r}^{m_1 \dots m_r}(k)]$. The first one is the sign relation denoted by *sign*. More precisely, let (F_1, f_1, O_1) and (F_2, f_2, O_2) be two forests. We write $(F_1, f_1, O_1) = \pm(F_2, f_2, O_2)$ if (F_1, f_1, O_1) differs from (F_2, f_2, O_2) only by reordering of the orientation set. The sign is the Koszul sign of permutation taking into account the degrees of the elements. The second relation is the Arnold's relation, denoted by *Arnold*, which is of the form



Arnold's relation

This relation is local, that is, all other vertices and edges are the same in each summand. Finally there is a differential in $\mathbb{Q}[\mathcal{F}_{s_1 \dots s_r}^{m_1 \dots m_r}(k)]$ defined as follows. For an edge $e \in E_F$, let $(F/e, f, O \setminus \{e\})$ be the forest obtained from (F, f, O) by contracting e . Recall that every vertex in F is labeled by some subset $A_i \subset A$. When an edge between two sets A_i and A_j is contracted, the new vertex is assigned the set $A_i \sqcup A_j$. Define a sign $(-1)^e \in \{-1, 1\}$ by $(-1)^e = (-1)^{\sum_{x < e} |x|}$. The differential of (F, f, O) is defined by

$$d(F, f, O) = \sum_{e \in E_F} (-1)^e (F/e, f, O \setminus \{e\})$$

The sequence $\left\{ \frac{\mathbb{Q}[\mathcal{F}_{s_1 \dots s_r}^{m_1 \dots m_r}(k)]}{\text{sign, Arnold}} \right\}_{k \geq 0}$ is equipped with a structure of right Ω -module defined as follows. Let $g: A' \rightarrow A$ be a surjective map from a finite set A' (of cardinal k') to A (of cardinal k). Define

$$g^*: \frac{\mathbb{Q}[\mathcal{F}_{s_1 \dots s_r}^{m_1 \dots m_r}(A)]}{\text{sign, Arnold}} \rightarrow \frac{\mathbb{Q}[\mathcal{F}_{s_1 \dots s_r}^{m_1 \dots m_r}(A')]}{\text{sign, Arnold}}$$

as the map sending a forest (F, f, O) to the forest $g^*(F, f, O)$ obtained by replacing each vertex of F labeled by a subset $A_i \subseteq A$ by a vertex labeled by $A'_i = g^{-1}(A_i)$. Notice that the orientation and the coloration of $g^*(F, f, O)$ remain unchanged. It is easy to see that the right Ω -module $CQ_{s_1 \dots s_r}^{m_1 \dots m_r}$ defined by (4.4) is exactly $\frac{\mathbb{Q}[\mathcal{F}_{s_1 \dots s_r}^{m_1 \dots m_r}]}{\text{sign, Arnold}}$. We reiterate, one just need to recall and apply the description of coLie in terms of spaces of trees modulo Arnold relations [39].

Now we will describe the Koszul dual $KQ_{s_1 \dots s_r}^{m_1 \dots m_r}$ of $Q_{s_1 \dots s_r}^{m_1 \dots m_r}$. It is defined as

$$KQ_{s_1 \dots s_r}^{m_1 \dots m_r} = Q_{s_1 \dots s_r}^{m_1 \dots m_r} \circ \text{coLie}[1].$$

So $CQ_{s_1 \dots s_r}^{m_1 \dots m_r}$, viewed as a right Com_+ module (we can view it like this because of (4.4) above), is freely generated by the symmetric sequence $KQ_{s_1 \dots s_r}^{m_1 \dots m_r}$. By definition we have an inclusion $KQ_{s_1 \dots s_r}^{m_1 \dots m_r} \subseteq CQ_{s_1 \dots s_r}^{m_1 \dots m_r}$, and generators of $KQ_{s_1 \dots s_r}^{m_1 \dots m_r}$ are forests (F, f, O) whose all vertices are singletons. Notice that for each k , $KQ_{s_1 \dots s_r}^{m_1 \dots m_r}(k)$ is concentrated in the single homological degree

$$\sum_{i=1}^r m_i s_i + \sum_{i=1}^r (k_i - s_i) = \sum_{i=1}^r (s_i(m_i - 1) + k_i) = \sum_{i=1}^r s_i(m_i - 1) + k,$$

where k_i is the number of vertices colored by i (so $k = k_1 + k_2 + \dots + k_r$).

The Koszul dual $KQ_{s_1 \dots s_r}^{m_1 \dots m_r}$ is isomorphic to a symmetric sequence related to the homology of configuration spaces as we will see in Proposition 4.2 below. Before stating this proposition we recall some notation. For a pointed topological space X , and for $p \geq 1$, let $\Delta^p X$ denote the "fat diagonal" in $X^{\wedge p} = \underbrace{X \wedge \dots \wedge X}_p$ (recall that the symbol " $\wedge$ " is that of the smash product). More precisely, we have

$$\Delta^p X = \bigcup_{1 \leq i \neq j \leq p} \{(x_1, \dots, x_p) \in X^{\wedge p} \mid x_i = x_j\}.$$

Proposition 4.2. For $s_1, \dots, s_r \geq 0$, and for $k \geq 0$ there is an isomorphism of symmetric sequences

$$KQ_{s_1 \dots s_r}^{m_1 \dots m_r}(\bullet) \cong \bigoplus_{\bullet = k_1 + \dots + k_r} \left(\text{Ind}_{\Sigma_{\bar{k}}}^{\Sigma_k} \bigotimes_{i=1}^r \tilde{H}_{(m_i-1)s_i+k_i}(S^{m_i k_i} / \Delta^{k_i} S^{m_i}, \mathbb{Q}) \right), \quad (4.6)$$

where $\Sigma_{\bar{k}} = \Sigma_{k_1} \times \dots \times \Sigma_{k_r}$.

Proof. The first thing to note is the following. For $1 \leq i \leq r$ one has

$$\tilde{H}_*(S^{m_i k_i} / \Delta^{k_i} S^{m_i}, \mathbb{Q}) = \overline{H}_*(C(k_i, \mathbb{R}^{m_i}), \mathbb{Q}), \quad (4.7)$$

where $\overline{H}_*(-)$ is the locally compact singular homology functor. The rest of the proof uses the Poincaré duality between $\overline{H}_*(C(k, \mathbb{R}^m), \mathbb{Q})$ and $\overline{H}^{m-k-*}(C(k, \mathbb{R}^m), \mathbb{Q})$, and also the fact that the latter cohomology admits a nice description in terms of forests (see [1, 8]). $\square$

Remark 4.3. There is a more natural topological way to prove (4.6). In [2, Section 11] one can find a general statement, saying that for any pointed space X , the Koszul dual of $\tilde{C}(X^{\wedge \bullet})$ is equivalent to $\tilde{C}_*(X^{\wedge \bullet} / \Delta^{\bullet} X)$. In our case $X = \vee_{i=1}^r S^{m_i}$, because of the formality the singular chains can be replaced by the homology:

$$\begin{aligned} K\tilde{H}_*((\vee_{i=1}^r S^{m_i})^{\wedge \bullet}, \mathbb{Q}) &\cong \tilde{H}_*((\vee_{i=1}^r S^{m_i})^{\wedge \bullet} / \Delta^{\bullet} (\vee_{i=1}^r S^{m_i}), \mathbb{Q}) \\ &\cong \bigoplus_{k_1 + \dots + k_r = \bullet} \text{Ind}_{\Sigma_{\bar{k}}}^{\Sigma_k} \tilde{H}_*(\wedge_{i=1}^r (S^{m_i k_i} / \Delta^{k_i} S^{m_i}), \mathbb{Q}), \end{aligned}$$

where as before $\Sigma_{\bar{k}} = \Sigma_{k_1} \times \dots \times \Sigma_{k_r}$.

4.2 Koszul complex computing the rational homology of $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$

In this subsection we describe a complex computing the rational homology of $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$.

Recall the following definition.

Definition 4.4. Let $V = \{V(n)\}_{n \geq 0}$ and $W = \{W(n)\}_{n \geq 0}$ be two symmetric sequences. Define a new symmetric sequence $V \hat{\otimes} W$ by

$$V \hat{\otimes} W(n) = \bigoplus_{p+q=n} \text{Ind}_{\Sigma_p \times \Sigma_q}^{\Sigma_n} V(p) \otimes W(q).$$

In practice we need to deal with multigraded vector spaces. Besides the usual homological degree, they will have the Hodge multi-grading $(s_1, \dots, s_r)$ and grading by complexity t . As usual when we take tensor product all the degrees get added.

Proposition 4.5. For $d > 2\max\{m_i \mid 1 \leq i \leq r\} + 1$, there is a quasi-isomorphism

$$\begin{aligned} C_*(\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \mathbb{Q}) &\simeq \left(\bigoplus_{k \geq 0} \text{hom}_{\Sigma_k} \left(\bigotimes_{1 \leq i \leq r} \overline{H}_*(C(\bullet, \mathbb{R}^{m_i}), \mathbb{Q})(k), \hat{H}_*(C(k, \mathbb{R}^d), \mathbb{Q}) \right), \partial \right) \\ &\simeq \left(\bigoplus_{k_1, \dots, k_r} \text{hom}_{\Sigma_{k_1} \times \dots \times \Sigma_{k_r}} \left(\bigotimes_{1 \leq i \leq r} \overline{H}_*(C(k_i, \mathbb{R}^{m_i}), \hat{H}_*(C(k_1 + \dots + k_r, \mathbb{R}^d), \mathbb{Q}) \right), \partial \right). \end{aligned} \quad (4.8)$$

The differential ∂ here is defined similarly to the $r = 1$ case [3, Subsection 5.2]. We won't describe it explicitly here. However, in next Subsection 4.3 we describe explicitly the dual complex computing the cohomology of the space of links.

Proof. It follows essentially from Proposition 4.1 and Proposition 4.2. The idea is to replace each term in the right hand side of (4.2) by $\text{Rmod}_{\Omega}(CQ_{s_1 \dots s_r}^{m_1 \dots m_r}, \widehat{H}_{t(d-1)}(C(\bullet, \mathbb{R}^d); \mathbb{Q}))$ (since, in Section 4.1, we have seen that $CQ_{s_1 \dots s_r}^{m_1 \dots m_r}$ is a cofibrant replacement of $Q_{s_1 \dots s_r}^{m_1 \dots m_r}$), and use the fact that $CQ_{s_1 \dots s_r}^{m_1 \dots m_r}$ is freely generated by $KQ_{s_1 \dots s_r}^{m_1 \dots m_r}$. The rest of the proof is very similar to [3, Section 5.2]. $\square$

4.3 Graph-complex computing the rational cohomology of $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$

In this subsection we will explicitly describe a graph-complex, denoted $HH^{m_1, \dots, m_r, d}$, computing the rational cohomology of $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$ and which is dual to (4.8). Notice that the graph-complex $HH^{m, d}$ was described in [3, Section 5.3]. In a similar way we will describe here $HH^{m_1, \dots, m_r, d}$. Our complex will be colored by the set $\{1, \dots, r\}$ of components of our links.

By taking the dual of (4.8), and recalling " $\widehat{\otimes}$ " from Definition 4.4, we have a quasi-isomorphism

$$C^*(\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \mathbb{Q}) \quad (4.9)$$

$$\simeq \left(\bigoplus_{k \geq 0} \bigoplus_{k=k_1+\dots+k_r} \left(\text{Ind}_{\Sigma_{k_1} \times \dots \times \Sigma_{k_r}}^{\Sigma_k} \overline{H}_{-*}C(k_1, \mathbb{R}^{m_1}) \otimes \dots \otimes \overline{H}_{-*}C(k_r, \mathbb{R}^{m_r}) \right) \otimes_{\Sigma_k} \widehat{H}^*C(k, \mathbb{R}^d), d \right) \quad (4.10)$$

$$\simeq \left(\bigoplus_{k \geq 0} \bigoplus_{k=k_1+\dots+k_r} \left(\overline{H}_{-*}C(k_1, \mathbb{R}^{m_1}) \otimes \dots \otimes \overline{H}_{-*}C(k_r, \mathbb{R}^{m_r}) \right) \otimes_{\Sigma_{k_1} \times \dots \times \Sigma_{k_r}} \widehat{H}^*C(k, \mathbb{R}^d), d \right) \quad (4.11)$$

By using the Poincaré duality, one has

$$\overline{H}_{-*}C(k_i, \mathbb{R}^{m_i}) \cong H^{*+m_i k_i}C(k_i, \mathbb{R}^{m_i}) \otimes (\text{sign}_{k_i})^{\otimes m_i}, 1 \leq i \leq r. \quad (4.12)$$

The equations (4.11) and (4.12) tell us that the cochain complex $C^*(\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \mathbb{Q})$ is quasi-isomorphic to a complex whose building blocks are the cohomology groups of configuration spaces, which are well known (see for example [1, 8]). Recall that $H^*(C(p, \mathbb{R}^n))$ is an algebra generated by elements $g_{ij}, 1 \leq i \neq j \leq n$, of degree $n-1$, modulo relations

$$g_{ij} = (-1)^n g_{ji}, g_{ij}^2 = 0, \text{ and the Arnold relation: } g_{ij}g_{jk} + g_{jk}g_{ki} + g_{ki}g_{ij} = 0.$$

This cohomology algebra admits a nice description in terms of certain spaces of forests. To every non-zero monomial, one can assign a forest putting an edge from vertex i to vertex j for each factor g_{ij} . Using this description, and using (4.12) and (4.11), we have the graph-complex $HH^{m_1, \dots, m_r, d}$ (computing $H^*(\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \mathbb{Q})$), which is described as follows. It is a graded vector space spanned by some set, denoted $\mathcal{G}$, of graphs, modulo certain relations. An element in $\mathcal{G}$ is a quadruple (G, f_G^v, f_G^e, O_G) , where

- $G = (V_G, E_G)$ is a graph, possibly disconnected, in which V_G is a finite set of non-labeled vertices (vertices are non-labeled because the last tensor product in (4.10) is taken over Σ_k), and E_G is a finite set of oriented edges;

- $f_G^v: V_G \rightarrow \{1, \dots, r\}$ is a function (possibly non-surjective) called a *coloration* of vertices, and $f_G^e: E_G \rightarrow \{0, 1, \dots, r\}$ is a similar *coloration* of edges. We say that elements in $(f_G^v)^{-1}(i)$ are vertices colored by i . Similarly for edges, elements in $(f_G^e)^{-1}(i)$, $1 \leq i \leq r$, are said *colored* (by i), while the elements in $(f_G^e)^{-1}(0)$ are said *non-colored* edges. The latter edges (corresponding to generators of $H^*(C(k, \mathbb{R}^d), \mathbb{Q})$) can connect any two vertices, while colored edges (corresponding to generators of $H^*(C(k_i, \mathbb{R}^{m_i}), \mathbb{Q})$) connect only two vertices having the same color (see Figure 7 in which $r = 2$).



Figure 7: A graph in $\mathcal{G}$

There are two restrictions on such graphs. If we remove all colored edges, the resulting graph is a forest in which each connected component has at least two vertices. On the other hand, if we remove all non-colored edges, the resulting graph is a forest with any type of connected components;

- O_G is a total order of the orientation set of G , where the latter one is defined in two equivalent ways. One switches from one description to another by applying the Poincaré duality (4.12). The first way naturally reminds the Bott-Taubes integration. The second way is algebraically more natural from the point of view of Koszul duality. In the first way, the orientation set is the union of the set of
 - non-colored oriented edges (considered as elements of degree $d - 1$),
 - colored oriented edges (an edge colored by i is of degree $m_i - 1$), and
 - vertices (a vertex colored by i is of degree $-m_i$).

In the second way, it is the union of the set of

- non-colored oriented edges (considered as elements of degree $d - 1$),
- colored non-oriented edges (of degree -1), and
- connected components with respect to colored edges (such a component, whose edges are colored by i , is of degree $-m_i$).

Let $\text{Vect}[\mathcal{G}]$ denote the vector space over $\mathbb{Q}$ spanned by $\mathcal{G}$, modulo the following relations.

- $G = \pm G'$ if G differs from G' only by the order of the orientation set. The sign $\pm$ is the Koszul sign of the corresponding permutation taking into account the degrees of the elements;
- $G = (-1)^d G'$ if G differs from G' only by the orientation of a non-colored edge;
- $G = (-1)^{m_i} G'$ if G differs from G' only by the orientation of an edge colored by i ;
- The Arnold relations with respect to both colored and non-colored edges.

Notice that the third relation is valid only when we consider the first definition of an orientation set. The differential, denoted δ , in $\text{Vect}[\mathcal{G}]$ is the sum of contractions of colored edges. In the second description of orientation sets, the sign is defined by pulling on the first place of the orientation set the colored edge which

is contracted and then removing it from this set. For the first description of orientation sets, the sign is slightly more complicated. We need to pull on respectively first and second places of the orientation set the contracted colored edge and the vertex, which is its source (to recall this edge is oriented), and then remove both of them.

Definition 4.6.

$$HH^{m_1, \dots, m_r, d} = (\text{Vect}[\mathcal{G}], \delta).$$

The following result is an immediate consequence of (4.10), (4.11), and (4.12).

Theorem 4.7. *For $d > 2\max\{m_i \mid 1 \leq i \leq r\} + 1$, we have an isomorphism*

$$H^*(HH^{m_1, \dots, m_r, d}) \cong H^*(\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \mathbb{Q}).$$

Remark 4.8. *Notice that disjoint union of graphs defines a product on $HH^{m_1, \dots, m_r, d}$. Moreover the latter complex is a free graded commutative algebra generated by its subcomplex $HH_\pi^{m_1, \dots, m_r, d}$ of connected graphs (with respect to both colored and non-colored edges). Therefore the homology of $HH_\pi^{m_1, \dots, m_r, d}$ is naturally isomorphic to the space dual to the rational homotopy of $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$.*

4.4 Koszul complex computing the rational homotopy of $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$

Similarly to Proposition 4.5, we get:

Proposition 4.9. *For $d > 2\max\{m_i \mid 1 \leq i \leq r\} + 1$, there is a quasi-isomorphism*

$$\begin{aligned} \pi_* \overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) \otimes \mathbb{Q} &\simeq \left(\bigoplus_{k \geq 0} \text{hom}_{\Sigma_k} \left(\widehat{\otimes}_{1 \leq i \leq r} \overline{H}_*(C(\bullet, \mathbb{R}^{m_i}), \mathbb{Q})(k), \widehat{\pi}_* C(k, \mathbb{R}^d) \otimes \mathbb{Q} \right), \partial \right) \\ &\simeq \left(\bigoplus_{k_1, \dots, k_r} \text{hom}_{\Sigma_{k_1} \times \dots \times \Sigma_{k_r}} \left(\bigotimes_{1 \leq i \leq r} \overline{H}_*(C(k_i, \mathbb{R}^{m_i}), \widehat{\pi}_* C(k_1 + \dots + k_r, \mathbb{R}^d) \otimes \mathbb{Q} \right), \partial \right). \end{aligned} \quad (4.13)$$

The differential ∂ here is defined similarly to the case of the homology complex. For the $r = 1$ case, see [3, Subsection 5.2].

Proof. Analogous to the proof of Proposition 4.5. □

4.5 Truncations of the Koszul complexes and the Goodwillie-Weiss tower

Contrary to the hairy graph-complexes, the Koszul complexes are well suited to determine rationally the Goodwillie-Weiss approximations for $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$.

Theorem 4.10. *For $d - \max\{m_i \mid 1 \leq i \leq r\} > 2$, one has*

$$\begin{aligned} T_n C_* (\overline{\text{Emb}}_c (\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \mathbb{Q}) &\simeq \left(\bigoplus_{0 \leq k \leq n} \text{hom}_{\Sigma_k} \left(\bigotimes_{1 \leq i \leq r} \overline{H}_*(C(\bullet, \mathbb{R}^{m_i}), \mathbb{Q})(k), \widehat{H}_*(C(k, \mathbb{R}^d), \mathbb{Q}) \right), \partial \right) \\ &\simeq \left(\bigoplus_{\substack{k_1, \dots, k_r \\ k_1 + \dots + k_r \leq n}} \text{hom}_{\Sigma_{k_1} \times \dots \times \Sigma_{k_r}} \left(\bigotimes_{1 \leq i \leq r} \overline{H}_*(C(k_i, \mathbb{R}^{m_i}), \widehat{H}_*(C(k_1 + \dots + k_r, \mathbb{R}^d), \mathbb{Q})) \right), \partial \right). \end{aligned} \quad (4.14)$$

Notice that the range is improved compared to Proposition 4.5, essentially because we do not need the convergence $T_\infty C_* (\overline{\text{Emb}}_c (\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \mathbb{Q}) \simeq C_* (\overline{\text{Emb}}_c (\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \mathbb{Q})$ (which does require $d > 2\max\{m_i \mid 1 \leq i \leq r\} + 1$).

Since the homotopy limit does not commute with singular chains functor, in general

$$C_* (T_n \overline{\text{Emb}}_c (\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \mathbb{Q}) \not\simeq T_n C_* (\overline{\text{Emb}}_c (\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \mathbb{Q}).$$

Thus the methods of this paper are not applicable to understand rationally $T_n \overline{\text{Emb}}_c (\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$ even in the range $d > 2\max\{m_i \mid 1 \leq i \leq r\} + 1$. However, we believe in the following:

Conjecture 4.11. *For $d - \max\{m_i \mid 1 \leq i \leq r\} > 2$, one has*

$$\begin{aligned} \pi_* T_n \overline{\text{Emb}}_c (\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) \otimes \mathbb{Q} &\simeq \left(\bigoplus_{0 \leq k \leq n} \text{hom}_{\Sigma_k} \left(\bigotimes_{1 \leq i \leq r} \overline{H}_*(C(\bullet, \mathbb{R}^{m_i}), \mathbb{Q})(k), \widehat{\pi}_* C(k, \mathbb{R}^d) \otimes \mathbb{Q} \right), \partial \right) \\ &\simeq \left(\bigoplus_{\substack{k_1, \dots, k_r \\ k_1 + \dots + k_r \leq n}} \text{hom}_{\Sigma_{k_1} \times \dots \times \Sigma_{k_r}} \left(\bigotimes_{1 \leq i \leq r} \overline{H}_*(C(k_i, \mathbb{R}^{m_i}), \mathbb{Q}), \widehat{\pi}_* C(k_1 + \dots + k_r, \mathbb{R}^d) \otimes \mathbb{Q} \right), \partial \right). \end{aligned} \quad (4.15)$$

This conjecture can be viewed as a refinement of Conjecture 3.1 and for $r = 1$ has also been shown in [16].

Proof of Theorem 4.10. Let Ω_n , respectively Γ_n , denote the full subcategory of Ω , respectively Γ , consisting of sets, respectively pointed sets, of cardinal $\leq n$, respectively $\leq n + 1$. The Pirashvili cross effect functor restricts to an equivalence of categories also for truncated modules:

$$\text{cr}: \text{Rmod}_{\Gamma_n} \longrightarrow \text{Rmod}_{\Omega_n}.$$

Similarly to (1.4), which holds for $d - \max\{m_i \mid 1 \leq i \leq r\} > 2$, one gets

$$T_n C_* (\overline{\text{Emb}}_c (\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \mathbb{Q}) \simeq \text{hRmod}_{\Gamma_n} (C_* ((\bigvee_{i=1}^r S^{m_i})^{\times \bullet}), H_*(C(\bullet, \mathbb{R}^d); \mathbb{Q})). \quad (4.16)$$

Now using the fact that truncation commutes with the cross effect and also the formality of the Ω -module $\widetilde{C}_*((\bigvee_{i=1}^r S^{m_i})^{\wedge \bullet}, \mathbb{Q})$, we get

$$\begin{aligned} T_n C_* (\overline{\text{Emb}}_c (\prod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \mathbb{Q}) &\cong \text{hRmod}_{\Omega_n} \left(\widetilde{H}_*((\bigvee_{i=1}^r S^{m_i})^{\wedge \bullet}, \mathbb{Q}), \widehat{H}_*(C(\bullet, \mathbb{R}^d), \mathbb{Q}) \right) \\ &\simeq \prod_{s_1, \dots, s_r, t} \text{hRmod}_{\Omega_n} \left(Q_{s_1 \dots s_r}^{m_1 \dots m_r}, \widehat{H}_{t(d-1)}(C(\bullet, \mathbb{R}^d), \mathbb{Q}) \right) \simeq \prod_{s_1, \dots, s_r, t} \text{hRmod}_{\Omega} \left(Q_{s_1 \dots s_r}^{m_1 \dots m_r}, tr_n \widehat{H}_{t(d-1)}(C(\bullet, \mathbb{R}^d), \mathbb{Q}) \right), \end{aligned} \quad (4.17)$$

where the functor $tr_n: \text{Rmod}_\Omega \rightarrow \text{Rmod}_\Omega$ is defined by

$$tr_n M(k) = \begin{cases} M(k), & k \leq n; \\ 0, & k > n. \end{cases}$$

The last equality in (4.17) follows from [45, Proposition 4.8].⁵

Now replacing $Q_{s_1 \dots s_r}^{m_1 \dots m_r}$ by $CQ_{s_1 \dots s_r}^{m_1 \dots m_r}$, we notice that only finitely many factors in (4.17) are non-zero, and the obtained complex is exactly the truncated (4.8). $\square$

Remark 4.12. *The space $\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d)$ can also be studied by the multivariable manifold calculus [33] similarly to the case of classical (all $m_i = 1$) string links considered in [34]. We expect that analogously to Theorem 4.10, one can show that for $d - \max\{m_i \mid 1 \leq i \leq r\} \geq 2$,*

$$T_{n_1 \dots n_r} C_*(\overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d), \mathbb{Q}) \simeq \left(\bigoplus_{k_i \leq n_i, i=1 \dots r} \text{hom}_{\Sigma_{k_1} \times \dots \times \Sigma_{k_r}} \left(\bigotimes_{1 \leq i \leq r} \overline{H}_*(C(k_i, \mathbb{R}^{m_i}), \hat{H}_*(C(k_1 + \dots + k_r, \mathbb{R}^d), \mathbb{Q})) \right), \partial \right). \quad (4.18)$$

We also conjecture that in the same range,

$$\pi_* T_{n_1 \dots n_r} \overline{\text{Emb}}_c(\coprod_{i=1}^r \mathbb{R}^{m_i}, \mathbb{R}^d) \otimes \mathbb{Q} \simeq \left(\bigoplus_{k_i \leq n_i, i=1 \dots r} \text{hom}_{\Sigma_{k_1} \times \dots \times \Sigma_{k_r}} \left(\bigotimes_{1 \leq i \leq r} \overline{H}_*(C(k_i, \mathbb{R}^{m_i}), \hat{\pi}_* C(k_1 + \dots + k_r, \mathbb{R}^d) \otimes \mathbb{Q}) \right), \partial \right). \quad (4.19)$$

References

- [1] V. I. Arnol'd, *The cohomology ring of the group of dyed braids*, Mat. Zametki 5 (1969), p. 227–231.
- [2] G. Arone and V. Turchin, *On the rational homology of high dimensional analogues of spaces of long knots*, Geom. Topol., vol. 18 (2014) 1261–1322.
- [3] G. Arone and V. Turchin, *Graph-complexes computing the rational homotopy of high dimensional analogues of spaces of long knots*, Ann. Inst. Fourier, Grenoble 65 (2015), no. 1, 1–62.
- [4] D. Bar-Natan, *Vassiliev homotopy string link invariants*. J. Knot Theory Ramifications 4 (1995), no. 1, 13–32.
- [5] P. Boavida de Brito and M. Weiss. *Spaces of smooth embeddings and configuration categories*. Preprint arXiv:1502.01640.
- [6] A. J. Casson, *Link cobordism and Milnor's invariant*, Bull. London Math. Soc. 7 (1975), 39–40.

⁵This also explains why the hairy graph-complexes can not be used to study the Goodwillie-Weiss approximations: the functor tr_n does not preserve injectivity.

- [7] S. Chmutov, S. Duzhin, and J. Mostovoy, *Introduction to Vassiliev knot invariants*. Cambridge University Press, Cambridge, 2012. xvi+504.
- [8] F. R. Cohen. The homology of C_{n+1} spaces. In F. Cohen, T. Lada, and J. May, *The homology of iterated loop spaces*, Springer-Verlag, Berlin, 1976, Lecture Notes in Mathematics, Vol. 533.
- [9] J. Conant, R. Schneiderman, and P. Teichner, *Tree homology and a conjecture of Levine*. Geom. Topol. 16 (2012), no. 1, 555–600.
- [10] D. Crowley, S. C. Ferry, and M. Skopenkov. *The rational classification of links of codimension > 2* . Forum Math. 26 (2014), no. 1, 239–269.
- [11] J. Ducoulombier and V. Turchin, *Delooping manifold calculus tower on a closed disc*. Paper to appear.
- [12] W. Dwyer and K. Hess, *Long knots and maps between operads*, Geom. Topol., vol. 16 (2012), no. 2, 919–955.
- [13] W. Dwyer and K. Hess, *A delooping of the space of string links*. Preprint arXiv:1501.00575.
- [14] W. Dwyer and K. Hess, *Delooping the space of long embeddings*. Paper to appear.
- [15] B. Fresse, *Koszul duality of operads and homology of partition posets*, in Homotopy theory: relations with algebraic geometry, group cohomology, and algebraic K -theory, Contemp. Math., vol. 346, Amer. Math. Soc., Providence, RI, 2004, p. 115–215.
- [16] B. Freese, V. Turchin, and T. Willwacher. *Mapping spaces of the E_n operads*. Paper to appear.
- [17] E. Getzler and M. Kapranov, *Cyclic operads and cyclic homology*. Geometry, topology, & physics, 167–201, Conf. Proc. Lecture Notes Geom. Topology, IV, Int. Press, Cambridge, MA, 1995.
- [18] E. Getzler and M. Kapranov, *Modular operads*. Compositio Math. 110 (1998), no. 1, 65–126.
- [19] T. Goodwillie and J. Klein, *Multiple disjunction for spaces of smooth embeddings*, J. Topol. 8 (2015), no. 3, 651–674.
- [20] T. G. Goodwillie and M. Weiss, *Embeddings from the point of view of immersion theory: Part II*, Geom. Topol. 3 (1999) 103–118.
- [21] N. Habegger and G. Masbaum, *The Kontsevich integral and Milnor’s invariants*. Topology 39 (2000), no. 6, 1253–1289.
- [22] N. Habegger and W. Pitsch, *Tree level algebra structures of perturbative invariants*, J. Knot Theory Ramifications 12 (2003), no. 3, 333–345.
- [23] A. Haefliger, *Knotted $(4k - 1)$ -spheres in $6k$ -space*, Ann. of Math. (2) 75 1962 452–466.
- [24] A. Haefliger, *Enlacements de sphères en codimension supérieure à 2*, Comm. Math. Helv. 41 (1966-67), 51-72 (in French).
- [25] V. Hinich and A. Vaintrob, *Cyclic operads and algebra of chord diagrams*. Selecta Math. (N.S.) 8 (2002), no. 2, 237–282.
- [26] M. Hovey, *Model categories*, Mathematical Surveys and Monographs 63, AMS 1999.

- [27] U. Koschorke, *A generalization of Milnor's μ -invariants to higher-dimensional link maps*, Topology 36 (1997), no. 2, 301–324.
- [28] P. Lambrechts and I. Volić, *Formality of the little N -disks operad* (2013), Memoirs of the American Mathematical Society, Volume 230, Number 1079.
- [29] J. Levine, *Addendum and correction to: "Homology cylinders: An enlargement of the mapping class group"*, Algebr. Geom. Topol. 2 (2002), 1197–1204.
- [30] J.-L. Loday and B. Vallette, *Algebraic operads*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 346, Springer, Heidelberg, 2012, xxiv+634 pages.
- [31] M. Markl, *Loop homotopy algebras in closed string field theory*. Comm. Math. Phys. 221 (2001), no. 2, 367–384.
- [32] B. Munson, *Derivatives of the identity and generalizations of Milnor's invariants*, J. Topol. 4 (2011), no. 2, 383–405.
- [33] B. Munson and I. Volić, *Multivariable manifold calculus of functors*, Forum Math. 24 (2012), no. 5, 1023–1066.
- [34] B. A. Munson and I. Volić, *Cosimplicial model for spaces of links*, J. of Homotopy Relat. Struct. 9 (2014), no. 2, 419–454.
- [35] T. Pirashvili, *Dold-Kan type theorem for Γ -groups*, Math. Ann. 318 (2000), no. 2, 277–298.
- [36] T. Pirashvili, *Hodge decomposition for higher order Hochschild homology*, Ann. Sci. Ecole Norm. Sup (4) 33 (2000), no. 2, 151–179.
- [37] K. Sakai, *Configuration space integrals for embedding spaces and the Haefliger invariant*, J. Knot Theory Ramifications 19 (2010), no. 12, 1597–1644.
- [38] K. Sakai and T. Watanabe, *1-loop graphs and configuration space integral for embedding spaces*, Math. Proc. Cambridge Philos. Soc. 152 (2012), no. 3, 497–533.
- [39] D. Sinha, *A pairing between graphs and trees*, preprint arXiv:math/0502547.
- [40] P. A. Songhafouo Tsopméné, *The rational homology of spaces of long links* (2013), Algebr. Geom. Topol. 16 (2016), no. 2, 757–782.
- [41] P. A. Songhafouo Tsopméné and Victor Turchin, *Euler characteristics for the Hodge splitting in the rational homology and homotopy of high dimensional string links*, to appear.
- [42] V. Turchin, *Hodge-type decomposition in the homology of long knots*, J. Topol. 3 (2010), no. 3, 487–534.
- [43] V. Turchin. *Delooping totalization of a multiplicative operad*. J. Homotopy Relat. Struct. 9 (2014), no. 2, 349–418.
- [44] V. Turchin and T. Willwacher. *Relative (non-)formality of the little cubes operads and the algebraic Cerf lemma*. Preprint arXiv:1409.0163.
- [45] V. Turchin and T. Willwacher. *Hochschild-Pirashvili homology on suspensions and representations of $\text{Out}(F_n)$* . Preprint arXiv:1507.08483.

- [46] M.Weiss, *Homology of spaces of smooth embeddings*, Q. J. Math. 55 (2004), no. 4, 499-504.
- [47] S. Whitehouse, *Gamma Homology of Commutative Algebras and Some Related Representations of the Symmetric Group*, PhD Thesis, Warwick University, 1994.

Institut de Recherche en Mathématique et Physique (IRMP)
Université Catholique de Louvain, Chemin du Cyclotron 2
1348 Louvain-la-Neuve, Belgique
E-mail address: arnaud.songhafouo@uclouvain.be

Department of Mathematics, Kansas State University
138 Cardwell Hall, Manhattan, KS 66506, USA
E-mail address: turchin@ksu.edu